IMPILO ® YEZIKELEMU WORM SYRUP

SCHEDUING STATUS: S0

PROPRIETARY NAME AND DOSAGE FORM:
IMPILO ® YEZIKELEMU WORM SYRUP (Syrup)

COMPOSITION: Each 5 ml contains:
Piperazine hydrate…….....625,0 mg
Preservative:
Sodium benzoate .....…….....0,1 % *m/v*
Excipients: caramel, citric acid monohydrate, sorbitol solution and Sweet Orange Flavour no.1.
Sugar free

CATEGORY AND CLASS: A 12 Anthelmintics

PHARMACOLOGICAL ACTION: Piperazine is an anthelmintic against roundworms (Ascaris) and threadworms (Enterolius). Piperazine produces reversible muscle paralysis at all stages of maturity except in larval forms. The worms are then expelled in the faeces by peristalsis.

INDICATIONS: For the elimination of roundworm and threadworm.

CONTRAINDICATIONS: IMPILO® YEZIKELEMU WORM SYRUP should not be taken by patients suffering from epilepsy, neurological disturbances, or impaired renal or hepatic function.

WARNINGS AND SPECIAL PRECAUTIONS:
Do not exceed the recommended dose unless prescribed by a doctor.
Contains sorbitol and may have a laxative effect.
Patients with the rare hereditary condition of sorbitol intolerance should not take Impilo Yezikelemu Worm Syrup.

HUMAN REPRODUCTION:
The safety of this preparation in pregnancy and lactation has not been established.

DOSAGE AND DIRECTIONS FOR USE:
Fasting before treatment is not necessary.
ROUNDWORM:
Dosage to be taken ONCE, preferably in the morning.
ADULTS AND CHILDREN OVER 12 YEARS:
A single dose of 6-7 medicine measurefuls
9-12 YEARS : 6 medicine measurefuls
6-8 YEARS : 5 medicine measurefuls
4-5 YEARS : 4 medicine measurefuls
1-3 YEARS : 2 medicine measurefuls
UNDER 1 YEAR: 1 ml / kg body mass
The dose may be repeated after 14 days.

THREADWORM: Treatment is for 7 days taken ONCE DAILY
ADULTS AND CHILDREN OVER 12 YEARS:
3 – 4 medicine measurefuls.
4 - 12 YEARS : 2 medicine measurefuls
2 - 3 YEARS: 1 medicine measureful
UNDER 2 YEARS: 0,36 – 0,6 ml / kg bodymass.
A second course after a seven day interval may be required. To avoid reinfection, it is advised that the entire household be treated.

SIDE EFFECTS:
Nausea, vomiting, diarrhoea, allergic reactions, abdominal pain, headache and urticaria may be experienced.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:
If large amount accumulate in the body due to overdosage or renal dysfunction, toxic effects may be experienced. Symptoms of overdosage include: drowsiness, dizziness, muscular inco-ordination or weakness, numbness, blurred vision, vertigo, confusion, tremor, convulsions, loss of reflexes and respiratory depression.
Toxic effects usually disappear when use is discontinued. The medicine may induce or exacerbate epileptic seizures in predisposed patients. Consult a doctor or the nearest hospital. Treatment is symptomatic and supportive.

IDENTIFICATION: A clear black syrup with a characteristic orange odour and a sweetish orange taste.

PRESENTATION: 25 ml Amber PVC bottle with white caps.

STORAGE INSTRUCTIONS: Store at or below 25 °C. Do not refrigerate. Protect from light. Keep the bottle tightly closed.
KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER: H/12/0881

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:
Avid Brands S.A. (Pty) Ltd
Suite 9, Hillcrest Office Park
2 Old Main Road, Hillcrest, 3610
www.avidbrands.com

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION:
23 May 2001

IMPILO ® YEZIKELEMU WORM STROOP

SKEDULERINGSTATUS: S0

EIENDOMSNAAM EN DOSEERWORM:
IMPILO ® YEZIKELEMU WORM SYRUP (Stroop)

SAMESTELLING: Elke 5 ml bevat:
Piperasienhidraat…….....625,0 mg
Preserveermiddel:
Natriumbensoaat .....……..0,1 % *m/v*
Onaktiewe bestanddele: karamel, sitroensuur monohidraat, sorbitol oplossing en Sweet Orange Flavour no. 1.
Suiker vry

KATEGORIE EN KLAS: A12 Teenhelmintiek middels.

FARMAKOLOGIESE WERKING: Piperasien is 'n anthelmintikum teen rondewurm (Ascaris) en haarwurm (Enterolius). Piperasien omkeerbare spierverlamming veroorsaak in rondewurm in alle stadiums van volwassenheid behalwe larvaalvorms. Die wurms word dan in die ontlasting deur dermbeweging uitgestoot.

INDIKASIES: Vir die verwydering van rondewurm en haarwurm.

KONTRAINDIKASIES: IMPILO® YEZIKELEMU WORM SYRUP moet nie deur pasiënte wat aan epilepsie, neurologiese-versteurings of verswakte nier – of lewerfunksie ly geneem word nie.

WAARSKUWINGS EN SPESIALE VOORSORGMAATREËLS:
Moet nie die aanbevole dosis oorskry tensy deur 'n geneesheer aanbeveel nie.
Bevat sorbitol en mag 'n lakserende effek hê.
Pasiënte met die seldsame oorerflike toestand van sorbitol intoleransie moet nie Impilo Yezikelemu Worm Syrup neem nie.

MENSLIKE REPRODUKSIE:
Die veiligheid van hierdie medisyne tydens swangerskap en borsvoeding, is nog nie vasgestel nie.

DOSIS EN GEBRUIKSAANWYSINGS:
Sonder kos bly voor behandeling is nie nodig nie.
RONDEWURM:
Alle dosisse moet EEN maal, in die oggend, geneem word.
VOLWASSENES EN KINDERS 12 JAAR EN OUER:
'n Enkel dosis van 6-7 medisynemate.
9-12 JAAR: 6 medisynemate
6-8 JAAR: 5 medisynemate
4-5 JAAR: 4 medisynemate
1-3 JAAR: 2 medisynemate
ONDER 1 JAAR: 1 ml / kg liggaams gewig.
Hierdie dosis kan na 14 dae herhaal word.

HAARWURM: Behandling is vir 7 aaneenlopende dae, EEN KEER PER DAG.
VOLWASSENES EN KINDERS 12 JAAR EN OUER:
3 - 4 medisynemate
4 - 12 JAAR: 2 medisynemate
2 - 3 JAAR: 1 medisynemate
ONDER 2 JAAR: 0,36 – 0,6 ml / kg liggaams gewig.
Herhaal behandeling na 'n een week tussentyd indien nodig. Om herinfeksie te vermy, is dit aanbeveel dat die hele huisgesin behandel word.

NEWE EFFEKTE:
Naarheid, vomering, diaree, allergiese reaksies, buikpyn, hoofpyn of netelroos mag ervaar word.

BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN:
Indien groot hoeveelhede in die liggaam ophoop, as gevolg van oordosering of nierdisfunksie, mag vergiftiging ervaar word. Simptome van oordosering sluit slaperigheid, duiseligheid, verwarring, 'n gebrek aan spierkoördinasie of spier swakheid, belemmerde visie, verwarring, siddering, verlies van reflekse en asemhalingsneerdrukking in. Vergiftigings simptome verdwyn gewoonlik wanneer gebruik beïndig word. Die medisyne mag epileptiesebeslagleggings in ontvanklike pasiënte veroorsaak of vererg. Raadpleeg 'n geneesheer of die naaste hospital. Behandeling is simptomaties en ondersteunend.

IDENTIFIKASIE: 'n helder Swart stroop met 'n lemoen geur en 'n soet lemoen smaak.

AANBIEDING: 25 ml Amber PVC bottels met wit proppe.

BERGINGSAANWYSINGS: Berg by of bende 25 °C.
Moenie verkoel nie. Beskerm teen lig. Hou die bottel dig toe.
HOU BUITE BEREIK VAN KINDERS.

REGISTRASIENOMMER: H/12/0881

NAAM EN BESIGHTIDSADRES VAN DIE HOUER VAN DIE REGISTRASIE SERTIFIKAAT:
Avid Brands S.A. (Edms) Bpk
Suite 9 Hillcrest Office Park
2 Old Main Straat, Hillcrest, 3610
www.avidbrands.com

DATUM VAN PUBLIKASIE VAN DIE PROFESSIONELE INLIGTING:
23 Mei 2001