1.3.1.1 PROFESSIONAL INFORMATION FOR MEDICINES FOR HUMAN USE

SCHEDULING STATUS S0

PROPRIETARY NAME AND DOSAGE FORM

LENNON GLYCERIN SUPPOSITORIES FOR INFANTS AND CHILDREN

COMPOSITION

Each suppository of LENNON GLYCERIN SUPPOSITORIES FOR INFANTS AND CHILDREN contains 1,12 g glycerin

Excipients:

Purified water, sodium carbonate anhydrous, sodium hydroxide, stearic acid

CATEGORY AND CLASS

A 11.6 Lubricants and faecal softeners

PHARMACOLOGICAL ACTION

Pharmacodynamic properties

Acts by its osmotic properties to attract water to the lumen, thereby softening the stool and promoting bowel movement.

INDICATIONS

For the relief of occasional constipation.

CONTRAINDICATIONS

- Sensitivity to the ingredients.
- Do not use unless the infant and child to be treated is in fact constipated.

WARNINGS AND SPECIAL PRECAUTIONS

FOR RECTAL USE ONLY.

This medicine should not be used in the presence of abdominal pain, nausea or vomiting.

Frequent or prolonged use of this preparation may result in dependence on laxatives and loss of normal bowel function.

If a sudden change in bowel habits is noticed that has persisted for a period greater than two weeks, consult a doctor before using the laxative.

No laxative should be used for longer than one week, except on the advice of a doctor.

Rectal bleeding or failure to have a bowel movement after use of a laxative may indicate a serious condition. Discontinue use and consult a doctor.


May cause rectal discomfort or a burning sensation (see SIDE EFFECTS).

Must be administered with caution to diabetic patients or to those at risk of dehydration.


DOSAGE AND DIRECTIONS FOR USE

Directions for removing suppository from wrapping

CUT ONE SUPPOSITORY OFF



AT THE SLIT PULL ONE SIDE OF THE FOIL FORWARD TO RELEASE SUPPOSITORY



Children 2 to under 6 years of age:

One to one and a half paediatric suppositories to be inserted rectally in a single daily dose, when necessary.

Children under 2 years of age:

One to one and a half paediatric suppositories to be inserted rectally in a single daily dose, when necessary.

Hold the wide base of the suppository in the fold of a cloth. Lubricate the pointed end with petroleum jelly or moisten with water and insert well up into the rectum of the child or infant to be treated. Even though complete solution is not necessary for action, the suppository should be held for five minutes to induce a motion. Acts within 15 to 30 minutes.

SIDE EFFECTS

Glycerin when used rectally may cause rectal discomfort or a burning sensation (see WARNINGS AND SPECIAL PRECAUTIONS).

Large doses of glycerin, when taken orally, may lead to serious toxic effects.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENTS

Symptoms

See SIDE EFFECTS and WARNINGS AND SPECIAL PRECAUTIONS.

Treatment

Treatment is symptomatic and supportive.

IDENTIFICATION

A smooth, clear to pale yellow, translucent torpedo shaped suppository.

PRESENTATION

12 suppositories are packed in an aluminium foil blister, inside bright side - outside matt side.

One strip is packed into an outer cardboard carton together with a leaflet.

STORAGE INSTRUCTIONS

Store at or below 25 °C.

Protect from moisture.

Keep in original packaging until required for use.

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER

E/11.6/1540

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF

REGISTRATION

PHARMACARE LIMITED

Healthcare Park

Woodlands Drive

Woodmead 2191

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION FOR MEDICINES FOR HUMAN USE

Date of registration: 23 April 1996

Date of the most recent amendment to the professional information as approved by the Authority: 23 April 1996

Botswana: B9315080 S3

Namibia: NS0 04/11.6/0033

Zimbabwe: P. 2017/16.5.2/5422

ZA_LEGLISUP_9604_00