IMPILO ® MILK OF MAGNESIA

SCHEDULING STATUS: S0

PROPRIETARY NAME AND DOSAGE FORM:
IMPILO ® MILK OF MAGNESIA (Suspension)

COMPOSITION:
Each 5 ml medicine measure contains:

Magnesium hydroxide	425 mg
Preservative: Sodium benzoate	0,1% *m/v*

Excipients: citric acid monohydrate and purified water.

Sugar free.

CATEGORY AND CLASS
A 11.4 Medicines acting on the gastro-intestinal tract.

PHARMACOLOGICAL ACTION:
Magnesium hydroxide has an acid neutralizing capacity and diminishes the activity of pepsin in gastric secretion. In large doses, it acts as a saline laxative by osmotic action.

INDICATIONS:
Where the use of an antacid is indicated. For the relief of occasional constipation.

CONTRAINDICATIONS:
Not to be taken by patients with impaired renal function, debility or dehydration, unless under the advice and supervision of a doctor. Must not be given to patients who are hypersensitive to any of the ingredients.

WARNINGS AND SPECIAL PRECAUTIONS:
Do not use this product of you have kidney disease, except under the advice and supervision of a doctor. Not to be taken by persons suffering from diarrhoea, abdominal pain, nausea or vomiting, unless directed by a doctor. If a sudden change in bowel habits is noticed, that persists over a period of 2 weeks, consult a doctor before using the laxative. Rectal bleeding or failure to have bowel movement after use of a laxative may indicate a serious condition. Discontinue use and consult a doctor.

INTERACTIONS:
This preparation may interfere with the absorption of other medicines if take concomitantly.

HUMAN REPRODUCTION:
The safety of this preparation in pregnancy and lactation has not been established.

DOSAGE AND DIRECTIONS FOR USE:
SHAKE BOTTLE BEFORE USE.
ANTACID:
Adults: 5 -10 ml (one or two medicine measures)
Children 3 -10 years: 2,5 ml (half a medicine measure)
NOT MORE THAN THREE TO FOUR TIMES PER 24 HOUR PERIOD.

LAXATIVE:
When used as a laxative, the medicine should be followed with sufficient water to prevent net loss of body water. Take as a single dose, or as directed by a doctor. This product generally produces bowel movement in half to six hours.
Adults: 20 - 40 ml (four to eight medicine measures)
Children 2 - 6 years: 5 -10 ml (one to two medicine measures)
6 -12 years: 10 - 20 ml (two to four medicine measures)
The dosages recommended above should not be exceeded in a 24 hour period. Do not use the maximum daily dosage of this product for more than one week except under the advice and supervision of a doctor.

SIDE EFFECTS:
When used as an antacid, the medicine may cause diarrhoea. The release of carbon dioxide may cause discomfort. With very large amounts or in patients with impaired renal function, hypermagnesaemia may develop, with the following symptoms: flushing of the skin, thirst, hypotension, drowsiness, confusion, loss of tendon reflexes, muscle weakness, respiratory depression, cardiac arrhythmias, coma and cardiac arrest.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:
See also Side Effects and Warnings and Special Precautions. Overdosage may be characterized by excessive purging. If symptoms of hypermagnesaemia are observed, namely flushing of the skin, thirst, hypertension due to peripheral vasodilation, drowsiness, confusion, loss of tendon reflexes, respiratory depression, cardiac arrhythmias, coma and cardiac arrest, consult a doctor. Treatment is symptomatic and supportive.

IDENTIFICATION:
A white suspension.

PRESENTATION:
100 ml PVC bottles with white caps.

STORAGE INSTRUSTIONS:
Store between 5 and 25 ºC. Do not freeze.
Keep bottle tightly closed.
KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER:
E/11.4/764

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:
Avid Brands S.A. (Pty) Ltd
Suite 9 Hillcrest Office Park
2 Old Main Road
Hillcrest
www.avidbrands.com

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION:
November 1994

IMPILO ® MILK OF MAGNESIA

SKEDULERINGSTATUS: S0

EIENDOMSNAAM EN DOSSERVORM:
IMPILO ® MILK OF MAGNESIA (Suspensie)

SAMESTELLING:
Elke 5 ml medisynemaatvol bevat:

Magnesiumhidroksied	425 mg
Preserveermiddel: Natriumbensoaat	0,1% *m/v*

Onaktiewe bestanddele: sitroensuur monohidraat en gesuiwerde water.

Suiker vry.

KATEGORIE EN KLAS
A 11.4 Middels met uitwerking op die maagdermkanaal.

FARMAKOLOGIESE AKSIE:
Magnesiumhidroksied het 'n suur-neutraliserende vermoë, en verminder die werking van pepsien in maagsekresie. In groter dosisse werk dit as n' lakseermiddel deur osmotiese werking.

INDIKASIES:
Waar die gebruik van 'n teensuurmiddel aangedui is. Vir die verlighting van toevallige harlywigheid.

KONTRA INDIKASIES:
Hierdie medisyne moet nie deur pasiënte wat ann verswakte nierfunksie, swakheid, of dehidrasie ly, geneem word, tensy volgens die raad en onder die toesig van 'n geneesheer nie. Moet nie deur pasiënte wie hipsosensitief is vir enige een van die bestanddele gebruik word nie.

WAARSKUWINGS EN SPESIALE VOORSORGMAATREËLS:
Moet nie hierdie medisyne neem indien u aan niersiekte ly, tensy volgens die raad en onder die toesig van 'n geneesheer nie. Moet nie deur mense wat ann diarree, buikpyn, naarheid of vomering ly, geneem word, tensy deur 'n geneesheer aanbeveel nie. Indien 'n skielike verandering in ingewandsgewoontes wat langer as twee weke voortduur waargeneem word, raadpleeg 'n geneesheer voor gebruik van hierdie lakseermiddel. Rektale bloeding of afwesgheid van 'n ingewande stoelgang na gebruik van 'n lakseermiddel kan 'n ernstige toestand anndui. Staak gebruik en raadpleeg 'n geneesheer.

INTERAKSIES:
Indien tegelykertyd geneem, mag hierdie perparaat die opneming van ander medisyne verhinder.

MENSLIKE REPRODUKSIE:
Die veiligheid van hierdie medisyne tydens swangerskap en borsvoeding, is nog nie vasgestel nie.

DOSIS EN GEBRUIKSAANWYSINGS:
SKUD DIE BOTTEL VOOR GEBRUIK.
TEENSUUR:
Volwassenes: 5 -10 ml (een tot twee medisynemate)
Kinders 3 -10 jaar: 2,5 ml ('n halwe medisynemaat)
NIE MEER AS DRIE TOT VIER KEER IN 'N 24 UUR TYDPERK.

LAKSEERMIDDEL:
Wanneer as 'n lakseermiddel gebruik, moet die medisune deur 'n voldoende hoeveelheid water gevlog word om netto verlies van liggaamswater te vermy. Neem as 'n enkele dosis, of soos deur 'n geneesheer aanbeveel. Oor die algemeen kan hierdie medisyne, tussen 'n half en ses ure, ingewande storlgang veroorsaak.
Volwassenes: 20 - 40 ml (vier tot agt medisynemate)
Kinders 2 - 6 jaar: 5 -10 ml (een tot twee medisynemate)
6 -12 jaar: 10 -12 ml (twee tot vier medisynemate)
Die aanbevole dosis moet nie binne 'n 24 uur tydperf oorskry nie. Moet nie die hoogste daaglikse dosis vir meer as een week neem tensy met die raad en onder die toesig van 'n geneesheer nie.

NEWE EFFEKTE:
Wanneer as 'n teensuur gebruik, mag die medisyne diarree veroorsaak. Die vrylating van koolsuurgas mag ongemak versoorsaak. Met baie groot hoeveelhede, of in pasiënte wat aan verswakte nierfuksie ly, mag hipermagnesemie, met die volgende simptime voorkom: blosing van die vel, dors, hipotensie, sluimerigheid, verwarring, verloor van seningreflekse, spierswakheid, asemhalingsneerdrukking, hartonreëlmatigheid, bewusteloosheid en hartstilstand.

BEKENDE SIMPTOME VAN OORDOSERING EN DIE BEHANDELING DAARVAN:
Verwys ook na Newe Effekte en Waarskuwings en Spesiale Voorsorgmaatreëls. Oordosering mag tot oordrewe purgasie lei. Indien simptome van hipermagnesemie naamlik, blosing van die vel, dors, hipotensie, sluimerigheid, verwarring, verloor van sening-reflekse, spierswakheid, asemhalingsneerdrukking,hartonreël-matigheid, bewusteloosheid en hartstilstand waargeneem word, raadpleeg 'n geneesheer. Die behandeling is simptomaties en ondersteunend.

IDENTIFIKASIE:
'n Wit suspensie.

AANBIEDING:
100 ml PVC bottels met wit proppe.

BERGINGSINSTRUKSIES:
Bêre tussen 5 en 25 ^{o}C. Moet nie bevries nie.
Hou bottel dig toe.
HOU BUITE BEREIK VAN KINDERS.

REGISTRASIENOMMER:
E/11.4/764

NAAM EN BESIGHTIDSADRES VAN DIE HOUER VAN DIE REGISTRASIE SERTIFIKAAT:
Avid Brands S.A. (Edms) Bpk
Suite 9 Hillcrest Office Park
2 Old Main Straat, Hillcrest
www.avidbrands.com

DATUM VAN PUBLIKASIE VAN DIE PROFESSIONELE INLIGTING:
November 1994