SCHEDULING STATUS:

S0

PROPRIETORY NAME:

L.C.C.®-G GUAIPHENESIN COUGH REMEDY - ORANGE FLAVOUR

L.C.C.®-G GUAIPHENESIN COUGH REMEDY – PEPPERMINT FLAVOUR

(AND DOSAGE FORM) (SYRUP)

COMPOSITION:

L.C.C.®-G GUAIPHENESIN COUGH REMEDY - ORANGE FLAVOUR:

Each 10 ml contains:

Guaifenesin	200 mg
Nipastat (Preservative)	0,02 % *m/v*

Excipients: saccharin sodium; sodium cyclamate; propylene glycol; levomenthol; glycerol; purified water; sorbitol 70 % solution; xanthan gum; Colour Yellow sunset; Flavour Orange Liquid.

Sugar and alcohol free.

Contains sweeteners: saccharin sodium 16 mg and sodium cyclamate 13 mg.

L.C.C.®-G GUAIPHENESIN COUGH REMEDY – PEPPERMINT FLAVOUR

Excipients: saccharin sodium; sodium cyclamate; propylene glycol; levomenthol; glycerol; purified water; sorbitol 70 % solution; xanthan gum; Caramel 48000; Peppermint oil.

Sugar and alcohol free

Contains sweeteners: saccharin sodium 16 mg and sodium cyclamate 13 mg.

PHARMACOLOGIGAL CLASSIFICATION:

A 10.1 Antitussives and expectorants.

PHARMACOLOGICAL ACTION:

Guaifenesin has expectorant properties.

INDICATIONS:

Alleviation of cough.

CONTRAINDICATIONS:

Hypersensitivity to any of the ingredients. Pregnancy. Porphyria.

WARNINGS AND SPECIAL PRECAUTIONS:

Guaifenesin should not be taken for persistent cough such as occurs with smoking, asthma, emphysema or where cough is accompanied by excessive secretions except under the advice and supervision of a doctor.

A persistent cough may be a sign of a serious condition. If cough persists for more than one week, tends to recur or is accompanied by high fever, skin rash or persistent headache, or sore throat is present with cough, consult a doctor.

Guaiphenesin should not be given to children under 2 years unless recommended by a doctor.

Patients with the rare hereditary condition of sorbitol intolerance should not take **L.C.C.®-G GUAIPHENESIN COUGH REMEDY.**

Effects on ability to drive and use machinery:

L.C.C.®-G GUAIPHENESIN COUGH REMEDY has no known effect on the ability to drive and to use machinery.

PREGNANCY AND LACTATION:

L.C.C.®-G GUAIPHENESIN COUGH REMEDY should not be used during pregnancy and lactation. (see Contraindications).

DOSAGE AND DIRECTIONS FOR USE:

Adults, adolescents and geriatrics

10 ml to 20 ml (2 to 4 medicine measures) every 4 hours.

Children:

6 to 12 years:

5 ml to 10 ml (1 to 2 medicine measures) every 4 hours.

3 to 6 years

2,5 ml to 5 ml (½ to 1 medicine measure) every 4 hours.

SIDE EFFECTS

Gastrointestinal discomfort (including diarrhoea, nausea, vomiting and stomach pains). Dizziness, headache, skin rash and urticaria occur less frequently.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

Very large doses can cause nausea and vomiting.

Treatment is symptomatic and supportive.

IDENTIFICATION:

ORANGE FLAVOUR: Orange, clear, viscous syrup. Odour of orange and menthol. Taste – sweet, of orange and menthol.

PEPPERMINT FLAVOUR: Brown, slightly turbid, viscous syrup. Odour of peppermint and menthol. Taste- sweet, of peppermint and menthol.

PRESENTATION:

50 ml, 75 ml and 100 ml in clear or amber PVC bottles.

STORAGE INSTRUCTIONS:

Store at or below 25 °C. Protect from freezing.

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBERS:

ORANGE FLAVOUR: 33/10.1/0057

Namibia: 04/10.1/1573 – NS0

PEPPERMINT FLAVOUR: 33/10.1/0121

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:

Adcock Ingram Limited

1 New Road,

Erand Gardens,

Midrand, 1685

Private Bag X69,

Bryanston, 2021.

www.adcock.com

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION:

April 1999