1.3.1.1 PROFESSIONAL INFORMATION FOR MEDICINES FOR HUMAN USE

SCHEDULING STATUS **S0**

PROPRIETARY NAME AND DOSAGE FORM

BORSTOL COUGH REMEDY (emulsion)

COMPOSITION

Each 10 ml emulsion of BORSTOL COUGH REMEDY contains:

Ammonium chloride	29 mg
Infusion senegae conc.	0,1666 ml
Infusion buchu conc.	0,0572 ml
Camphor	0,2135 ml
Tinct. cinchona co. sine serpentaria	0,4453 ml
Balsam sulphuris	0,3125 ml
Turpentine Oil	0,3125 ml

Excipients:

Arachis oil, castor oil, chloroform, ethanol 96 %, ether, flavour peppermint oil, flavour tincture capsicum, linseed oil, menthol levo, purified water, saccharin sodium, sucrose refined white H1, tincture lavandulae co

Contains alcohol: Ethyl alcohol 11,0 % *v/v*

Contains sugar: Sucrose 724 mg

Contains sweetener: Saccharin sodium 7,0 mg

CATEGORY AND CLASS

A 10.1 Antitussives and expectorants

PHARMACOLOGICAL ACTION

Expectorant and demulcent.

INDICATIONS

BORSTOL COUGH REMEDY is indicated for the symptomatic relief of general and chronic troublesome coughs and hoarseness often associated with colds, influenza and bronchitis.

CONTRAINDICATIONS

Not known.

WARNINGS AND SPECIAL PRECAUTIONS

Excipients:

BORSTOL COUGH REMEDY contains sucrose which may have an effect on the glycaemic control of patients with diabetes mellitus.

Patients with rare hereditary conditions such as fructose intolerance, glucose-galactose mal-absorption or sucrase-isomaltase insufficiency should not take BORSTOL COUGH REMEDY.

INTERACTIONS

Not known.

HUMAN REPRODUCTION

Not known.

DOSAGE AND DIRECTIONS FOR USE

Shake the bottle well.

Adults: Take 5 ml to 10 ml every 2, 3 or 4 hours.

SIDE EFFECTS

Not known.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT

In case of an overdose, consult your physician.

IDENTIFICATION

Brown emulsion which separates into an upper oily brown layer and a lower brown aqueous layer on standing. On shaking the brown layer becomes lighter but more opaque.

PRESENTATION

50 ml, 100 ml or 200 ml are packed into an oval glass bottle and sealed with a black aluminium roll on cap that is internally lacquered and lined with an expanded polyethylene liner. The bottle is placed in an outer cardboard carton together with a leaflet.

Not all packs and pack sizes are necessarily marketed.

STORAGE INSTRUCTIONS

Store at or below 30 °C.

Keep in original packaging until required for use.

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER

G907 (Act 101/1965)

NAME AND BUSINESS ADDRESS OF THE APPLICANT

PHARMACARE LIMITED

Healthcare Park

Woodlands Drive

Woodmead 2191

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION FOR MEDICINES FOR HUMAN USE

Date of registration: Old Medicine

Date of the most recent amendment to the professional information as approved by the Authority: 02 November 1982

ZA_BORSCEML_8211_01