FLIXOTIDE

SCHEDULING STATUS:

S3

PROPRIETARY NAME AND DOSAGE FORM:

FLIXOTIDE 50 INHALER CFC-FREE

FLIXOTIDE 125 INHALER CFC-FREE

FLIXOTIDE 250 INHALER CFC-FREE

Pressurised metered-dose inhaler

COMPOSITION:

FLIXOTIDE 50 INHALER CFC-FREE delivers 50 micrograms of fluticasone propionate per actuation.

FLIXOTIDE 125 INHALER CFC-FREE delivers 125 micrograms of fluticasone propionate per actuation.

FLIXOTIDE 250 INHALER CFC-FREE delivers 250 micrograms of fluticasone propionate per actuation.

Excipients: hydroxyfluoroalkane 134a, 1,1,1,2-tetrafluoroethane (HFA 134a).

PHARMACOLOGICAL CLASSIFICATION:

A 21.5.1 Corticosteroids and analogues.

PHARMACOLOGICAL ACTION:

Pharmacodynamic properties:

Fluticasone propionate *in vitro* has a potent glucocorticoid anti-inflammatory action.

Pharmacokinetic properties:

Following oral administration 87 % - 100 % of the dose is excreted in the faeces, up to 75 % as parent compound depending on the dose. There is a non-active major metabolite. Following intravenous administration there is rapid plasma clearance suggestive of extensive hepatic extraction. The plasma elimination half-life is approximately 3 hours. The volume of distribution is approximately 250 litres.

INDICATIONS:

Prophylactic management of atopic asthma in adults and children of 6 years and older.

CONTRA-INDICATIONS:

FLIXOTIDE is contra-indicated in patients with a history of allergy to any of its components.

WARNINGS AND SPECIAL PRECAUTIONS:

Treatment with FLIXOTIDE should not be stopped abruptly as adrenal insufficiency may be precipitated in this way.

Systemic corticosteroid effects may occur in patients on fluticasone treatment.

Patients transferred from other inhaled steroids or oral steroids remain at risk of impaired adrenal reserve for a considerable time after transferring to inhaled fluticasone propionate.

Increasing use of short-acting inhaled beta-2 agonists to control symptoms indicates deterioration of asthma control. Under these conditions, the patient should be reassessed.

Sudden and progressive deterioration in asthma control is potentially life-threatening and may have several causes. Consideration should be given to increasing corticosteroid dosage if not caused by otherwise treatable causes of deterioration.

Severe asthma requires regular medical assessment as death may occur.

Patients with severe asthma may require high dose inhaled (see dosage instructions) or oral corticosteroid therapy. Sudden worsening of symptoms may require increased corticosteroid dosage which should be administered under urgent medical supervision.

Patients weaned off oral steroids whose adrenocortical function is still impaired should carry a steroid warning card indicating that they may need supplementary systemic steroid during periods of stress, e.g. worsening asthma attacks, chest infections, major intercurrent illness, surgery, trauma, etc.

In rare cases inhaled therapy may unmask underlying eosinophilic conditions (e.g. Churg Strauss syndrome). These cases have usually been associated with reduction or withdrawal of oral corticosteroid therapy. A direct causal relationship has not been established.

Similarly replacement of systemic steroid treatment with inhaled therapy may unmask allergies such as allergic rhinitis or eczema previously controlled by the systemic drug. These allergies should be symptomatically treated with antihistamine and/or topical preparations, including topical steroids.

Systemic effects may occur with any inhaled corticosteroid, particularly at high doses prescribed for long periods; these effects are much less likely to occur than with oral corticosteroids. Possible systemic effects include adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataract and glaucoma. It is important, therefore, that the dose of inhaled corticosteroid is titrated to the lowest dose at which effective control is maintained.

It is recommended that the height of children receiving prolonged treatment with inhaled corticosteroid is regularly monitored.

Patient's inhaler technique should be checked to make sure that inhaler actuation is synchronised with inspiration to ensure optimum delivery of the drug to the lungs.

Patients in a medical or surgical emergency, who require high doses of inhaled steroids and/or intermittent treatment with oral steroids, are at risk of impaired adrenal reserve.

The extent of the adrenal impairment may require specialist advice before elective procedures. The possibility of residual impaired adrenal response should always be borne in mind in emergency and elective situations likely to produce stress and appropriate corticosteroid treatment must be considered.

In children taking recommended doses of inhaled fluticasone propionate adrenal function and adrenal reserve usually remain within the normal range. However, the possible effects of previous or intermittent treatment with oral steroids should not be discounted.

Lack of response or severe exacerbations of asthma should be treated by increasing the dose of inhaled fluticasone propionate or by giving a systemic steroid and/or an antibiotic if there is an infection.

Special care is necessary in patients with active or quiescent pulmonary tuberculosis.

Patients on corticosteroid therapy may have adrenocortical suppression.

Effects on ability to drive and use machines:

FLIXOTIDE is unlikely to produce an effect.

INTERACTIONS:

No specific interaction studies have been performed.

PREGNANCY AND LACTATION:

Safety during pregnancy and lactation has not been established.

Corticosteroids have been shown to be teratogenic in animals. As these agents are absorbed when inhaled, teratogenicity following inhalation cannot be excluded.

DOSAGE AND DIRECTIONS FOR USE:

Shake before use.

FLIXOTIDE INHALER CFC-FREE is administered by the inhaled route only.

Patients should be made aware of the prophylactic nature of therapy with inhaled fluticasone propionate and that it should be taken regularly even when they are asymptomatic. The onset of therapeutic effect is 4 to 7 days. FLIXOTIDE INHALER CFC-FREE is not for relief in acute attacks but for routine long term management.

Patients will require a fast- and short-acting inhaled bronchodilator to relieve acute asthmatic symptoms.

The dosage of FLIXOTIDE INHALER CFC-FREE should be adjusted according to the individual response.

If patients find that relief with short-acting bronchodilator treatment becomes less effective or they need more inhalations than usual, medical attention must be sought.

Adults and children over 16 years of age:

100 to 1 000 micrograms twice daily.

Patients should be given a starting dose of inhaled fluticasone propionate which is appropriate for the severity of their disease. The dose may be adjusted until control is achieved or reduced to the minimum effective dose, according to the individual response.

Children over 6 years of age:

50 to 100 micrograms twice daily.

The dose may be adjusted until control is achieved and should be reduced to the minimum effective dose according to the individual response.

Special patient groups:

There is no need to adjust the dose in elderly patients.

For the transfer of patients being treated with oral corticosteroids:

- Patients who have been treated with systemic steroids for long periods of time or at a high dose may have adrenocortical suppression. With these patients adrenocortical function should be monitored regularly and their dose of systemic steroid reduced cautiously.
- After approximately a week, gradual withdrawal of the systemic steroid may be commenced. Decrements in dosages should be appropriate to the level of maintenance systemic steroid, and introduced at not less than weekly intervals. For maintenance doses of prednisolone (or equivalent) of 10 mg daily or less, the decrements in dose should not be greater than 1 mg per day, at not less than weekly intervals. For maintenance doses of prednisolone in excess of 10 mg daily, it may be appropriate to employ cautiously, larger decrements in dose at weekly intervals. In some patients on oral corticosteroids the dose reduction or replacement with inhaled corticosteroids may not be possible.
- Some patients feel unwell in a non-specific way during the withdrawal phase despite maintenance or even improvement of the respiratory function. They should be encouraged to persevere with inhaled fluticasone propionate and to continue withdrawal of systemic steroid, unless there are objective signs of adrenal insufficiency.

SIDE EFFECTS:

Candidiasis of the mouth and throat (thrush) may occur. Such patients may find it helpful to rinse out their mouth with water after using the inhaler. Symptomatic candidiasis can be treated with topical anti-fungal therapy whilst still continuing with the FLIXOTIDE.

Inhaled fluticasone propionate may cause hoarseness. It may be helpful to rinse out the mouth with water immediately after inhalation.

Paradoxical bronchospasm may occur with an immediate increase in wheezing after dosing. This should be treated immediately with a fast-acting inhaled bronchodilator. FLIXOTIDE should be discontinued immediately, the patient assessed, and if necessary alternative therapy instituted.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

Acute - Inhalation of the drug at dosages in excess of those recommended may lead to temporary suppression of adrenal function. This does not necessitate emergency action being taken. In these patients treatment with fluticasone propionate by inhalation should be continued at a dose sufficient to control asthma; adrenal function recovers in a few days and can be verified by measuring plasma cortisol.

Chronic - Use of inhaled fluticasone propionate in daily doses in excess of 2 milligrams over prolonged periods may lead to some degree of adrenal suppression. Monitoring of adrenal reserve may be indicated. Treatment with inhaled fluticasone propionate should be continued at a dose sufficient to control asthma.

IDENTIFICATION:

Metered-dose aerosol inhaler containing a white to off-white suspension.

PRESENTATION:

Pressurised metered-dose inhaler comprising an aluminium alloy can fitted with a metering valve.

FLIXOTIDE 50 INHALER CFC-FREE delivers 50 micrograms of fluticasone propionate per actuation.

FLIXOTIDE 125 INHALER CFC-FREE delivers 125 micrograms of fluticasone propionate per actuation.

FLIXOTIDE 250 INHALER CFC-FREE delivers 250 micrograms of fluticasone propionate per actuation.

Each canister of FLIXOTIDE 125 INHALER CFC-FREE and FLIXOTIDE 250 INHALER CFC-FREE provides 60 or 120 actuations. Each canister of FLIXOTIDE 50 INHALER CFC-FREE provides 120 actuations.

STORAGE INSTRUCTIONS:

Fluticasone propionate inhalers should be stored at or below 30 °C.

Protect from heat, frost and direct sunlight.

The canister should not be punctured, broken or burnt even when apparently empty.

Keep out of reach of children.

REGISTRATION NUMBER:

FLIXOTIDE 50 INHALER CFC-FREE: 35/21.5.1/0377

FLIXOTIDE 125 INHALER CFC-FREE: 32/21.5.1/0082

FLIXOTIDE 250 INHALER CFC-FREE: 32/21.5.1/0083

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:

GlaxoSmithKline South Africa (Pty) Ltd

39 Hawkins Avenue

Epping Industria 1, 7460

DATE OF PUBLICATION OF THE PACKAGE INSERT:

Date of registration:

FLIXOTIDE 50 INHALER CFC-FREE: 06 August 2002

FLIXOTIDE 125 INHALER CFC-FREE: 12 November 1999

FLIXOTIDE 250 INHALER CFC-FREE: 12 November 1999

Date of most recent revision: 05 August 2002

Date compliant with Regulation 11: 30 October 2015

GDS-7

MANUFACTURER:

Glaxo Wellcome Production

Zone Industrielle No. 2, 23 Rue Lavoisier, 27 000

Evreux, France

Botswana:
Flixotide 125 CFC Free Inhaler - Reg No BOT1001688A/B S2
Flixotide 250 CFC Free Inhaler - Reg No BOT9900429 S2
Malawi:
Flixotide 50 CFC Free Inhaler – Reg No PMPB/PL270/156 POM
Flixotide 125 CFC Free Inhaler – Reg No PMPB/PL270/73 POM
Flixotide 250 CFC Free Inhaler – Reg No PMPB/PL270/74 POM
Namibia:
Flixotide 50 CFC Free Inhaler - Reg No 04/21.5.1/0857 NS2
Flixotide 125 CFC Free Inhaler - Reg No 04/21.5.1/0856 NS2
Flixotide 250 CFC Free Inhaler - Reg No 04/21.5.1/0855 NS2

Zambia:
Flixotide 50 CFC Free Inhaler – Reg No 179/036 POM
Flixotide 125 CFC Free Inhaler – Reg No 179/035 POM
Flixotide 250 CFC Free Inhaler – Reg No 179/034 POM
Zimbabwe:
Flixotide 50 CFC Free Inhaler - Reg No 99/22.1.3/3470 PP
Flixotide 125 CFC Free Inhaler – Reg No 2015/22.1.3/5061 PP
Flixotide 250 CFC Free Inhaler – Reg No 2015/22.1.3/5062 PP

Trade marks are owned by or licensed to the GSK group of companies.

© 2017 GSK group of companies or its licensor.