SCHEDULING STATUS: **S1**

PROPRIETARY NAME AND DOSAGE FORM:

Iliadin® Kids Nose Drops

Iliadin® Kids Nose Spray

Nose Drops and Metered Nose Spray

COMPOSITION:

Each 1 ml of the solution contains:

0,25 mg/0,025% *m/v* Oxymetazoline Hydrochloride

0,005 % *m/v* Benzalkonium Chloride as preservative

PHARMACOLOGICAL CLASSIFICATION:

A.16.1 - Nasal Decongestants

PHARMACOLOGICAL ACTION:

Iliadin® Kids Nose Drops/Spray is a topical vasoconstrictor, which exerts a decongestant action on mucous membranes of the nasopharyngeal cavity.

INDICATIONS:

Iliadin® Kids Nose Drops/Spray is indicated for the relief of nasal congestion due to the common cold, sinusitis, allergic rhinitis and hay-fever.

Indicated as adjunctive treatment in middle ear infection.

CONTRAINDICATIONS:

Hypersensitivity to any of the ingredients.

WARNINGS AND SPECIAL PRECAUTIONS:

Iliadin® Kids Nose Drops/Spray should be used with care in patients with cardiac disease,

hypertension or hyperthyroidism.

Iliadin® Kids Nose Drops/Spray should not be used for a period longer than five days.

Prolonged use may cause rebound congestion and drug-induced rhinitis.

It is not advisable to use **Iliadin® Kids Nose Drops/Spray** more frequently than recommended.

INTERACTIONS:

None Known.

PREGNANCY AND LACTATION:

Iliadin® Kids Nose Drops/Spray is indicated for paediatric use only.

DOSAGE AND DIRECTIONS FOR USE:

Iliadin® Kids Nose Drops/Spray Nose Drops and Metered Nose Spray are indicated for children up to the age of six years.

Metered Nose Spray:

Two to three sprays into each nostril 2 – 3 times a day. See diagram for directions for use.

Nose Drops:

Two to three drops into each nostril 2 – 3 times a day. See diagram for directions for use.

SIDE EFFECTS:

Iliadin® Kids Nose Drops/Spray may cause local stinging or burning, sneezing and dryness of the mouth and throat.

If unexpected side effects appear, consult your doctor immediately.

KNOWN SYMPTOMS OF OVER DOSAGE AND PARTICULARS OF ITS TREATMENT:

See **SIDE EFFECTS.**

Local gastric irritation and possible systemic sympathomimetic effects may occur, e.g. sweating, tachycardia, drowsiness possibly leading to coma, convulsions and circulatory collapse.

Treatment is symptomatic and supportive.

IDENTIFICATION:

Clear, colourless solution, with a maximum of a faint odour.

PRESENTATION:

10 ml Metered Nose Spray (bottle).

10 ml Nose Drops (bottle).

STORAGE INSTRUCTIONS:

Store at or below 25 ˚C.

Protect from light.

Keep out of reach of children.

REGISTRATION NUMBER:

V/16.1/296.

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:

P&G South African Trading (Pty) Ltd.

12th Floor, Alice Lane Towers, 15 Alice Lane

Sandton, 2196

South Africa

DATE OF PUBLICATION OF THIS PACKAGE INSERT:

8 July 1988

[1] = Company Registration Number: 1989/004241/07

®= Registered Trademark

Namibia	NS1	04/16.1/0584

Botswana	S3	B9306415 (Nose Drops) B9306420 (Metered Nose Spray)

Iliadin® Kids Nose Drops/Spray

Directions for use

Metered Nose Spray

For best results keep both the head and spray bottle in an upright position. Remove cap by pulling and insert the nozzle of the spray loosely into the nostril with the middle and forefinger around the bottom of the nozzle and the thumb on the base of the bottle. Depress the spray mechanism (as shown in the diagram) thus producing a fine mist. Sniff in to ensure an even distribution of the spray. Withdraw from nostril and release the spray mechanism. Repeat for the other nostril.



Nose Drops

Fill glass dropper unit by placing the open end into the liquid and depressing the rubber teat.
Tilt the head back and instil 2- 3 drops of the liquid in each nostril.

