SCHEDULING STATUS: **S0**

PROPRIETARY NAME: **ANTIPEOL OINTMENT**
(AND DOSAGE FORM) **(OINTMENT)**

COMPOSITION:

Each 10 mg contains:

Zinc oxide	2 mg
Salicylic acid	0,01 mg
Ichthammol	0,093 mg

Contains urea

PHARMACOLOGICAL CLASSIFICATION:

A 13.8 Dermatological preparations. Keratolytics

PHARMACOLOGICAL ACTION:

Astringent, bacteriostatic, fungicidal and keratolytic actions.

INDICATIONS:

Chronic ulcers, dandruff, eczema, psoriasis, parasitic skin diseases and chronic skin diseases.

CONTRAINDICATIONS:

Hypersensitivity to any of the components.

WARNING:

For external use only.

DOSAGE AND DIRECTIONS FOR USE:

Apply the ointment first around and then on the affected area. Renew applications frequently after removing the old ointment with clean cotton wool or lint.

SIDE EFFECTS AND SPECIAL PRECAUTIONS:

Dermatitis may occur. Slight irritation to the skin. Should be used with care on large areas of skin. Symptoms of acute systemic salicylate poisoning have been reported after application to large areas of the body.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

As under "**SIDE EFFECTS**". If accidentally swallowed contact your doctor immediately or take the patient to the nearest hospital.

Treatment: Cease application and treat symptomatically.

IDENTIFICATION:

Smooth, beige coloured, unctuous ointment with a lavender odour.

PRESENTATION:

18 g, 37 g and 93 g aluminium ointment tubes.

STORAGE INSTRUCTIONS:

Store in a cool place below 25 °C in an airtight container.

KEEP OUT OF THE REACH OF CHILDREN.

REFERENCE NUMBER:

G2200 (Act 101/1965)

NAME AND BUSINESS ADDRESS OF THE APPLICANT:

Tiger Consumer Brands
2 Purlin Street
Isando
1600

DATE OF APPLICATION OF THIS PACKAGE INSERT:

6 December 1974

SKEDULERINGSTATUS: **S0**

EIENDOMSNAAM: **ANTIPEOL OINTMENT**
(EN DOSEERVORM) **(SALF)**

SAMESTELLING:

Eike 10 mg bevat:	
Sinkoksied	2 mg
Salisielsuur	0,01 mg
lgtammol	0,093 mg
Bevat ureum	

FARMAKOLOGIESE KLASSIFIKASIE:

A 13.8 Velpreparate. Keratolitika

FARMAKOLOGIESE WERKING:

Saamtrekkende, bakterie- en swamdodende en keratolitiese werkings.

INDIKASIES:

Kroniese rousere, skilfers, ekseem, psoriase, parasitiese velaandoenings en kroniese velsiektes.

KONTRA-INDIKASIES:

Oorgevoeligheid vir enige van die bestanddele.

WAARSKUWING:

Slegs vir uitwendige gebruik.

DOSIS EN GEBRUIKSAANWYSINGS:

Smeer die salf eers om en dan oor die aangetasde deel. Herhaal die aanwendings gereeld nadat die ou salf met skoon watte of verbandlinne verwyder is.

NEWE-EFFEKTE EN SPESIALE VOORSORGMAATREËLS:

Velontsteking kan voorkom. Ligte prikkeling op die vel. Behoort versigtig op groot veloppervlaktes gebruik te word. Simptome van akute sistemiese salisilaatvergiftiging is gemeld na aanwending oor groot liggaamsoppervlaktes.

BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN:

Soos onder “**NEWE-EFFEKTE**”.

lndien van die salf per ongeluk ingesluk word raadpleeg onmiddellik u geneesheer of neem die pasiënt na die naaste hospitaal.

Behandeling: Staak aanwending en behandel simptomaties.

IDENTIFIKASIE:

Gladde, roomkleurige, vetterige salf met 'n laventelreuk.

AANBIEDING:

18 g, 37 g en 93 g aluminium salfbuise.

BERGINGSAANWYSINGS:

Bêre in 'n koel plek onder 25 °C in 'n lugdigte houer.

HOU BUITE BEREIK VAN KINDERS.

VERWYSINGSNOMMER:

G2200 (Wet 101/1965)

NAAM EN BESIGHEIDSADRES VAN DIE APPLIKANT:

Tiger Consumer Brands
2 Purlin Street
Isando
Kempton Park
1600

DATUM VAN PUBLIKASIE VAN HIERDIE VOUBILJET:

6 Desember 1974