Patient Information Leaflet

SCHEDULING STATUS:

S4

PROPRIETARY NAME, STRENGTH AND PHARMACEUTICAL FORM:

FOXAIR 50/100 Accuhaler Powder for inhalation

FOXAIR 50/250 Accuhaler Powder for inhalation

FOXAIR 50/500 Accuhaler Powder for inhalation

Salmeterol 50 µg and fluticasone propionate 100 µg, 250 µg or 500 µg

Read all of this leaflet carefully before you start using FOXAIR Accuhaler.

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor or your pharmacist.
- FOXAIR has been prescribed for you or your child personally and should not be shared with other people. It may harm them, even if their symptoms are the same as yours.

WHAT FOXAIR ACCUHALER CONTAINS:

Each FOXAIR Accuhaler blister contains 50 micrograms of the active ingredient salmeterol (as xinafoate). Each blister also contains 100 or 250 or 500 micrograms of the active ingredient microfine fluticasone propionate.

Contains sugar (lactose up to 12,5 mg per blister) as excipient.

WHAT FOXAIR ACCUHALER IS USED FOR:

Your doctor has chosen this medicine to suit you and your condition. FOXAIR Accuhaler is used to help with asthma in people who need prevention treatment.

BEFORE YOU USE FOXAIR ACCUHALER:

Do not use FOXAIR Accuhaler:

- if you are allergic to salmeterol xinafoate
- if you are allergic to fluticasone propionate
- if you are allergic to lactose or milk proteins**.**

Take special care with FOXAIR ACCUHALER:

Special care should be taken with FOXAIR:

- if you ever had thrush in your mouth
- if you are being or have ever been treated for tuberculosis (TB)
- if you are having treatment for a thyroid condition
- if you are you having treatment for high blood pressure or a heart problem

Pregnancy and Breastfeeding:

If you are pregnant or breastfeeding your baby, please consult your doctor, pharmacist or other healthcare professional for advice before using FOXAIR.

Important information about some of the ingredients of FOXAIR Accuhaler:

Each blister of powder contains lactose/fructose. Patients with the rare hereditary conditions of lactose/fructose or galactose malabsorption should not take FOXAIR Accuhaler. Lactose may have an effect on the control of your blood sugar if you have diabetes mellitus.

Using other medicines with FOXAIR Accuhaler:

Always tell your healthcare professional if you are taking other medicines. (This includes complementary or traditional medicines.)

You must not take FOXAIR Accuhaler if you are receiving a medicine called ritonavir, without first consulting your doctor. Make sure your doctor knows what other medicines you are taking, including those that you have bought yourself.

HOW TO USE FOXAIR ACCUHALER:

DO NOT use this medicine to treat a sudden attack of breathlessness as it will not help you. You will need a different type of medicine, which you must not confuse with FOXAIR Accuhaler. You should always have your fast-acting 'rescue' medicine available. It is very important that you use it regularly every day. Do not stop treatment even if you feel better unless told to do so by your doctor. Do not change your dose unless told to by your doctor. If you are breathless or wheezing more often than normal, tell your doctor.

The medicine in the FOXAIR Accuhaler should be inhaled into your lungs. You will be able to taste the powder on your tongue if you have used the Accuhaler correctly. If you have any difficulties or do not understand the instructions ask your doctor or pharmacist. FOXAIR Accuhaler must only be breathed in through the mouth. Take your medicine as your doctor has told you. The label on it will usually tell you how much to take and how often. If you are not sure ask your doctor or pharmacist.

The usual dose is:

Asthma:

Adults and adolescents 12 years and older, as prescribed by your doctor:

One inhalation (FOXAIR 50/100 Accuhaler) twice daily or

One inhalation (FOXAIR 50/250 Accuhaler) twice daily or

One inhalation (FOXAIR 50/500 Accuhaler) twice daily.

Children 4 years and older:

One inhalation (FOXAIR 50/100 Accuhaler) twice daily.

COPD:


Adults: One inhalation (FOXAIR 50/250 Accuhaler) to one inhalation (FOXAIR 50/500 Accuhaler) twice daily.

Instructions for use of your FOXAIR Accuhaler:


Use and Handling:

The Accuhaler releases a powder which is inhaled into the lungs. A dose indicator on the Accuhaler indicates the number of doses left.

Closed: When you take your Accuhaler out of its box, it will be in the closed position.



Opened: A new Accuhaler contains 60 doses of your medicine. The dose indicator tells you how many doses are left.



This Accuhaler contains 60 individually protected doses of your medicine, in powder form. Each dose is accurately measured and hygienically protected. It requires no maintenance - and no refilling. The dose indicator on top of your Accuhaler tells you how many doses are left. Numbers

5 to 0 will appear in RED, to warn you when there are only a few doses left. The Accuhaler is easy to use. When you need a dose, just follow the four simple steps illustrated


1. Open.
2. Slide.
3. Inhale.
4. Close.

How your Accuhaler works:

Sliding the lever of your Accuhaler opens a small hole in the mouthpiece and unwraps a dose, ready for you to inhale it. When you close the Accuhaler, the lever automatically moves back to its original position, ready for your next dose when you need it. The outer case protects your Accuhaler when it is not in use.


1. **Open - How to use the Accuhaler.**

   To open your Accuhaler, hold the outer case in one hand and put the thumb of your other hand on the thumbgrip. Push your thumb away from you as far as it will go.



2. **Slide.**

   Hold your Accuhaler with the mouthpiece towards you. Slide the lever away from you, as far as it will go - until it clicks. Your Accuhaler is now ready to use. Every time the lever is pushed back, a dose is made available for inhaling. This is shown by the dose counter. Do not play with the lever as this releases doses which will be wasted.



3. Inhale.

Before you start to inhale the dose, read through this section carefully.

Hold the Accuhaler away from your mouth.

Breathe out as far as is comfortable. Remember - never breathe into your Accuhaler.


Put the mouthpiece to your lips.

Breathe in steadily and deeply - through the Accuhaler, not through your nose.

Remove the Accuhaler from your mouth.

Hold your breath for about 10 seconds, or for as long as is comfortable.


Breathe out slowly.



4. Close.

To close your Accuhaler, put your thumb in the thumbgrip, and slide the thumbgrip back towards you, as far as it will go.

When you close the Accuhaler, it clicks shut. The lever automatically returns to its original position and is reset. Your Accuhaler is now ready for you to use again.



If you have been instructed to take two inhalations you must close the Accuhaler and repeat stages 1 to 4.

Remember:

Keep your Accuhaler dry.

Keep it closed when not in use.

Never breathe into your Accuhaler.

Only slide the lever when you are ready to take a dose.

Do not exceed the stated dose. Keep out of reach of children.

After using your Accuhaler:

If your breathing or wheezing gets worse straight after using your Accuhaler or inhaler, stop using it immediately and tell your doctor as soon as possible.

If you use more FOXAIR Accuhaler than you should:

If you accidentally take a larger dose than recommended you may notice that your heart is beating faster than usual and that you feel shaky. Other symptoms you may experience include headache, muscle weakness and aching joints. Tell your doctor as soon as possible if you accidentally take a larger dose than you were told to.

If you missed a dose of FOXAIR Accuhaler:

If you miss a dose just take the next dose when it is due.

POSSIBLE SIDE EFFECTS:

FOXAIR can have side effects.

Not all side effects reported for FOXAIR are included in this leaflet. Should your general health worsen or if you experience any untoward effects while using FOXAIR, please consult your doctor, pharmacist or other healthcare professional for advise.

Some people can be allergic to medicines. If you have any of the following symptoms soon after taking FOXAIR Accuhaler, STOP taking this medicine and tell your doctor immediately.

- Sudden wheeziness and chest pain or tightness.
- Swelling of eyelids, face, lips, tongue or throat.
- Lumpy skin rash or "hives" anywhere on the body.

The above are all very serious side effects. If you have them, you may have had a serious allergic reaction to FOXAIR. You may need urgent medical attention, or hospitalisation.

Some people very occasionally find that their mouth, throat or tongue becomes sore or that their voice becomes hoarse after inhaling this medicine. Gargling with water and spitting it out immediately after each dose may help. Tell your doctor but do not stop treatment unless told to do so. Also tell your doctor if you have any of the following symptoms: headache, muscle cramps, pains in joints, skin rash or trembling, increase in pulse rate.

It is possible that some patients, particularly those taking higher doses of this type of medication, may very rarely suffer from the following side effects:

Rounded face, loss of bone density, eye problems and slowing of growth in children. If you have any other symptoms that you do not understand, tell your doctor or pharmacist.

STORING AND DISPOSING OF FOXAIR ACCUHALER:

Store in a dry place at or below 30 °C.

Store all medicines out of reach of children.

Do not use after the expiry date stated on your label or carton.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

PRESENTATION OF FOXAIR ACCUHALER:

A single Accuhaler is packed into a carton.

FOXAIR Accuhaler is a two-tone purple, circular moulded plastic device containing a foil strip with 60 regularly spaced blisters.

The FOXAIR 50/100 Accuhaler device is labelled with pink lettering.

The FOXAIR 50/250 Accuhaler device is labelled with purple lettering.

The FOXAIR 50/500 Accuhaler device is labelled with black lettering.

The device contains a dose counter which shows the number of doses remaining (60 to 1). To show when the last five doses have been reached, the number appears in red.

IDENTIFICATION OF FOXAIR ACCUHALER:

Each blister of FOXAIR 50/100, 50/250 and 50/500 Accuhaler contains a white powder.

REGISTRATION NUMBER:

FOXAIR 50/100 Accuhaler – 42/21.5.4/0581

FOXAIR 50/250 Accuhaler – 42/21.5.4/0582

FOXAIR 50/500 Accuhaler – 42/21.5.4/0583

NAME AND ADDRESS OF THE REGISTRATION HOLDER:

GlaxoSmithKline South Africa (Pty) Ltd

39 Hawkins Avenue

Epping Industria 1, 7460

DATE OF PUBLICATION:

Date of registration: 4 June 2010

Date of implementation of Regulation 12: 10 December 2015

GDS-10

Trade marks are owned by or licensed to the GSK group of companies.

© 2018 GSK group of companies or its licensor.

