CEZLIN CAPSULES
P1538046

CEZLIN CAPSULES
P1538046

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS S4

CEZLIN (150 mg) Capsules
Clindamycin hydrochloride
Contains sugar: 53,76 mg lactose monohydrate

Read all of this leaflet carefully before you start taking CEZLIN
- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist, nurse or other healthcare provider.
- CEZLIN has been prescribed for you personally and you should not share your CEZLIN with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet
1. What CEZLIN is and what it is used for
2. What you need to know before you use CEZLIN
3. How to use CEZLIN
4. Possible side effects
5. How to store CEZLIN
6. Contents of the pack and other information

1. What CEZLIN is and what it is used for
CEZLIN belong to a group of medicines called antibiotics. It is used to treat serious bacterial infections.
CEZLIN can be prescribed to you if you have any of the following:
- Upper respiratory tract infections such as inflammation of the lining of bronchial tubes, bronchial tubes (lung infection).
- Lower respiratory tract infections such as pain or irritation in the throat and inflammation of the nasal cavities.
- Skin and soft tissue infections, infected wounds, and dental infections.
- Inflammation of bone or bone marrow infections.

2. What you need to know before you take CEZLIN
Do not take CEZLIN
- If you are hypersensitive (allergic) to clindamycin, lincomycin or doxorubicin, or any of the other ingredients of CEZLIN (listed in Section 6).
- If you are suffering from diarrheal states or gastro-intestinal disease, particularly those with a history of colitis.
- If you are pregnant or breastfeeding.

Warnings and precautions
Take special care with CEZLIN
- Tell your doctor immediately if you have hypersensitivity skin reactions like fever, unexplained widespread skin pain, a red or purple skin rash that spreads or blisters on your skin and mucous membranes of your mouth, nose, eyes and genitals. CEZLIN should be discontinued and appropriate therapy should be initiated by your doctor.
- Tell your doctor immediately if you have diarrhoea or usually get diarrhoea when you take antibiotics or have ever suffered from problems with your stomach or intestines. If you develop severe or prolonged or bloody diarrhoea during or after using CEZLIN tell your doctor immediately since it may be necessary to stop the treatment. This may be a sign of bowel inflammation (pseudomembranous colitis) which can occur following treatment with antibiotics up to several weeks after discontinuation of treatment.
- If you have to take CEZLIN for a long time, your doctor may arrange regular liver, kidney and blood tests.
- If you suffer from problems with your kidneys or liver
- If you have rare hereditary problems of galactose intolerance or total lactase deficiency or glucose-galactose malabsorption you should not take this CEZLIN.

Other medicines and CEZLIN
Always tell your healthcare professional if you are taking any other medicine (this includes complementary or traditional medicines).
Do not take CEZLIN if
- If you are taking or using any nerve- and muscle pain blocking medicines.
- If you are taking or using any medicines related to any bacterial infections such as macrolides, ketolides, streptogramins, linezolid, and chloramphenicol.
- If you are taking or using a blood thinner or anti-blood clotting medicine such as warfarin.

Pregnancy, breastfeeding and fertility
If you are pregnant or breastfeeding your baby or think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare professional for advice before CEZLIN is administered to you.

Pregnancy
Safety and efficacy in pregnancy have not been established for CEZLIN.
CEZLIN is contraindicated in pregnancy There are no adequate and well-controlled studies in pregnant women during the first trimester of pregnancy.

Breastfeeding
You should not use CEZLIN if you are breastfeeding. Safety and efficacy in lactation have not been established. CEZLIN is excreted in the breast milk.

Fertility
CEZLIN do not have any effects on fertility or mating ability.

Driving and using machines
CEZLIN has no or negligible influence on the ability to drive and use machines.
It is not always possible to predict to what extent CEZLIN may interfere with your daily activities. You should ensure that you do not engage in the above activities until you are aware of the measure to which CEZLIN affects you.

CEZLIN contains lactose
Lactose is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this CEZLIN.

3. How to take CEZLIN
Do not share medicines prescribed for you with any other person. Always take CEZLIN exactly as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure.

Adults and elderly patients
The recommended dose is 150 mg - 450 mg (1-3 capsules) every six hours, depending on the severity of your infection. The capsules should be taken with a full glass of water.

Children
The recommended dose in children is between 3 and 6 mg per kg of body weight every six hours, depending on the severity of the infection. Your doctor will work out the number of capsules that your child should have.

Your doctor will tell you how long your treatment with CEZLIN will last. Do not stop treatment early. If you have the impression that the effect of CEZLIN is too strong or weak, tell your doctor or pharmacist.

Taking CEZLIN with food and drink
CEZLIN can be taken with food and water.

If you take more CEZLIN than you should
In the event of over dosage, consult your doctor or pharmacist without delay. If neither is available, contact the nearest hospital or poison control centre.

If you forget to take CEZLIN
It is important that you continue to take CEZLIN according to the instructions on the label and that you do not miss any doses. However, if you do miss a dose, take it as soon as you remember and take the next dose at the normal time. Do not take a double dose to make up for a forgotten dose.

Effects if CEZLIN is stopped
You must not stop taking CEZLIN unless your doctor tells you to. If you have any further questions on the use of CEZLIN, ask your doctor or pharmacist.

4. Possible side effects
CEZLIN can have side effects.
Not all side effects reported for CEZLIN are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking CEZLIN, please consult your doctor, pharmacist or other healthcare professional for advice.

If any of the following happens, stop taking CEZLIN and tell your doctor immediately or go to the casualty department at your nearest hospital:
- inflammation in the colon, loose stools, and abdominal pain, hypersensitivity, SJS (severe illness with ulceration of the mouth, lips and skin), TEN (severe illness and sloughing of the skin), AGEP (skin reactions), DRESS (a rash, involvement of internal organs, hematologic abnormalities, and systemic illness)
- severe, persistent or bloody diarrhoea (which may be associated with stomach pain or fever). This is a common side effect which may occur after treatment with antibiotics and can be a sign of serious bowel inflammation (pseudomembranous colitis)
- signs of a severe allergic reaction such as sudden wheeziness, difficulty in breathing, swelling of eyelids, face or lips, throat or tongue, rash or itching (especially affecting the whole body)
- a widespread rash with blistering and peeling of large areas of skin, particularly around the mouth, nose, eyes or genitals, known as Stevens-Johnson syndrome, or toxic epidermal necrolysis
- a rare skin eruption that is characterised by the rapid appearance of areas of red skin studded with small pustules (small blisters filled with white/yellow fluid) (Acute Generalised Exanthematous Pustulosis (AGEP))
- skin rash, which may blister, and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge - erythema multiforme)
- widespread red skin rash with small pus-containing blisters (bullous exfoliative dermatitis)
- fever, swollen lymph nodes or skin rash, these may be symptoms of a condition known as DRESS (Drug Reaction with Eosinophilia and Systemic Symptoms).

These are all very serious side effects. If you have them, you may have had a serious reaction to CEZLIN. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:
- Yellowing of the skin and eyes, dark urine, and tiredness which may be symptoms of liver problems.

These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:
Frequent side effects
- Vomiting
- Nausea (feeling sick)
- Abdominal cramps or stomach pains
- Rash maculo-papular (presence of flat and papules rashes)
- Urticaria itchy rash or swellings

Less frequent side effects
- *Clostridium difficile* colitis (mild diarrhoea to fatal colitis)
- vaginal infection
- Agranulocytosis (deficiency of granulocytes in the blood)
- Neutropenia (a low level of neutrophils)
- Eosinophilia (n increase in the number of eosinophils in the blood)
- Leukopenia (a reduction in the number of leucocytes) blood cells that fight for infection in the blood
- Thrombocytopenia (a reduction in the number of platelets in the blood)
- Dysgeusia (taste disturbance)
- Oesophageal ulcer
- Oesophagitis (inflammation of the lining of the oesophagus)
- Pruritus (itchy skin)
- rash morbilliform (skin rashes)

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects
If you get side effects, talk to your doctor, pharmacist or nurse.
This includes any possible side effects not listed in this leaflet. You can also report side effects via https://www.sahpra.org.za/documents/864221b6.04_ARF1_Jul16_v4.pdf.
By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store CEZLIN
STORE ALL MEDICINES OUT OF REACH OF CHILDREN.
- Store at or below 25 °C.
- Keep in original packaging until required for use.
- Do not use after the expiry date stated on the label.
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information
What CEZLIN contains
The active substance in CEZLIN is clindamycin hydrochloride USP equivalent to clindamycin 150 mg.

The other ingredients are
Lactose monohydrate
Corn starch
Talc
Magnesium stearate
Empty hard gelatin capsule shell (Light blue opaque / Light green transparent, Size '1')

What CEZLIN looks like and contents of the pack
Light blue opaque / Light green transparent size '1' hard gelatin capsule filled with white to off white powder.

HDPE Container Pack (20's)
White opaque round 40 ml HDPE container with 33 mm neck finish closed with 33 mm white opaque polypropylene RS closure with wad having induction sealing liner. Each HDPE container also contains 1 gm of silica gel sachet. HDPE containers are further packed in pre-printed cartons with a leaflet.

HDPE Container Pack (100's)
White opaque round 120 ml HDPE container with 38 mm neck finish closed with 38 mm white opaque polypropylene CR closure with wad having induction sealing liner. Each HDPE container also contains 1 gm of silica gel sachet. HDPE containers are further packed in pre-printed cartons with a leaflet.

Name, business address and telephone number of the holder of the certificate of registration
Novagen Pharma (Pty) Ltd
Office 2, 100 Sovereign Drive,
Route 21 Corporate Park,
Nellmapius Drive, Irene, 0157, Pretoria, South Africa
Tel. No.: +27 012 345 3175
Email address: info@novagenpharma.co.za

Registration number
52/20.1.1/0028

This leaflet was last revised in
12 October 2021

The Professional information for CEZLIN (150 mg) capsules is available on the Novagen Pharma website: http://www.novagenpharma.co.za/products/antibiotics/

Marketed by:
United Pharma Marketing (Pty) Ltd

2408A158

PASIËNTINLIGTINGSPAMFLET

SKEDULERINGSTATUS S4

CEZLIN (150 mg) Kapsules
Klindamisienhidrochloried
Bevat suiker : 53,76 mg laktosemonohidraat

Lees die hele pamflet noukeurig deur voordat jy CEZLIN begin neem.
- Bewaar hierdie pamflet. Jy sal dit dalk weer moet lees.
- Indien jy verdere vrae het, vra asseblief jou dokter of apteker, verpleegster of enige ander gesondheidsorgwerker.
- CEZLIN is persoonlik vir jou voorgeskryf en jy moenie jou medisyne met enige ander mense deel nie. Dit kan hulle benadeel, selfs al is hul simptome dieselfde as joune.

Wat is in hierdie pamflet
1. Wat CEZLIN is en waarvoor dit gebruik word
2. Wat jy moet weet voordat jy CEZLIN neem
3. Hoe om CEZLIN te neem
4. Moontlike newe-effekte
5. Hoe om CEZLIN te berg
6. Inhoud van die verpakking en ander inligting

1. Wat CEZLIN is en waarvoor dit gebruik word
CEZLIN behoort aan 'n groep medisyne wat antibiotika genoem word. Dit word gebruik om ernstige bakteriële infeksies te behandel.
CEZLIN kan aan jou voorgeskryf word as jy enige van die volgende het:
- Boonste lugweginfeksies soos inflammasie van die voering van bronigiale buise (longinfeksie).
- Laer lugweginfeksies soos pyn of irritasie in die keel en ontsteking van die neusholtes.
- Vel- en sagteweefselinfeksies, besmette wonde en tandheelkundige infeksies.
- Ontsteking van been- of beenmurginfeksies.

2. Wat jy moet weet voordat jy CEZLIN neem
Moenie CEZLIN neem nie
- As jy hipersensitief (allergies) is vir klindamisien, linkomisien of doksorubisien, of enige van die ander bestanddele van CEZLIN (gelys in Afdeling 6).
- As jy aan diarree of gastro-intestinale siektes ly, veral dié met 'n geskiedenis van kolitis.
- As jy swanger is of borsvoed.

Waarskuwings en voorsorgmaatreëls
Wees veral versigtig met CEZLIN
- Vertel jou dokter dadelik as jy hipersensitiewe velreaksies soos koors, onverklaarbare wydverspreide velpyn, 'n rooi of pers veluitslag wat versprei of blase op jou vel en slymvliese van jou mond, neus, oë en geslagsdele het. CEZLIN moet gestaak word en toepaslike terapie moet deur jou dokter begin word.
- Vertel jou dokter dadelik as jy diarree het of gewoonlik diarree kry wanneer jy antibiotika neem of al ooit aan probleme met jou maag of ingewande gely het. As jy ernstige of langdurige of bloederige diarree ontwikkel tydens of na die gebruik van CEZLIN, vertel jou dokter dadelik aangesien dit nodig mag wees om die behandeling te staak. Dit kan 'n teken wees van dermontsteking (pseudomembraneuse kolitis) wat kan voorkom na behandeling met antibiotika tot 'n paar weke na staking van behandeling.
- As jy CEZLIN vir 'n lang tyd moet neem, kan jou dokter reël vir gereelde lewer-, nier- en bloedtoetse.
- As jy probleme met jou niere of lewer het.
- As jy seldsame oorerflike probleme van galaktose-intoleransie of totale laktasetekort of glukose-galaktose-wanabsorpsie het, moet jy nie CEZLIN neem nie.

Ander medisyne en CEZLIN
Vertel altyd jou gesondheidsorgwerker as jy enige ander medisyne neem (dit sluit komplementêre of tradisionele medisyne in).
Moenie CEZLIN neem as
- As jy medikasie neem of gebruik wat senuwee- en spierpyn blokkeer.
- As jy enige medisyne neem of gebruik wat verband hou met enige bakteriële infeksies soos makroliede, ketoliede, streptogramiene, linezolid, en chloramfenikol.
- As jy 'n bloedverdunner of medisyne teen bloedstolling soos warfarin neem of gebruik.

Swangerskap, borsvoeding en vrugbaarheid
As jy swanger is of jou baba borsvoed of dink jy is dalk swanger of beplan om 'n baba te hê, raadpleeg asseblief jou dokter, apteker of ander gesondheidswerker vir raad voordat CEZLIN aan jou toegedien word.

Swangerskap
Veiligheid en doeltreffendheid tydens swangerskap is nie vir CEZLIN vasgestel nie.
CEZLIN is teenaangedui tydens swangerskap. Daar is geen voldoende en goed-gekontroleerde studies in swanger vroue gedurende die eerste trimester van swangerskap nie.

Borsvoeding
Jy moenie CEZLIN gebruik as jy borsvoed nie. Veiligheid en doeltreffendheid in laktasie is nie vasgestel nie. CEZLIN word in die borsmelk uitgeskei.

Vrugbaarheid
CEZLIN het geen uitwerking op vrugbaarheid of intimiteit nie.

Bestuur en gebruik van masjiene
CEZLIN het geen of 'n weglaatbare invloed op die vermoë om te bestuur en masjiene te gebruik nie.
Dit is nie altyd moontlik om te voorspel tot watter mate CEZLIN met jou daaglikse aktiwiteite kan inmeng nie. Jy moet verseker dat jy nie by die bogenoemde aktiwiteite betrokke raak voordat jy bewus is van die mate waarbe CEZLIN jou raak nie.

CEZLIN bevat laktose
Laktose is 'n soort suiker. As jou dokter jou vertel het dat jy 'n onverdraagsaamheid teenoor sommige suikers het, kontak jou dokter voordat jy hierdie CEZLIN neem.

3. Hoe om CEZLIN te neem
Moenie medisyne wat vir jou voorgeskryf is met enige ander persoon deel nie.
Neem altyd CEZLIN presies soos jou dokter vir jou voorgeskryf het. Jy moet jou dokter of apteker raadpleeg as jy onseker is.

Volwassenes en bejaarde pasiënte
Die aanbevole dosis is 150 mg - 450 mg (1-3 kapsules) elke ses uur, afhangende van die erns van jou infeksie. Die kapsules moet met 'n vol glas water geneem word.

Kinders
Die aanbevole dosis by kinders is tussen 3 en 6 mg per kg liggaamsgewig elke ses uur, afhangende van die erns van die infeksie. Jou dokter sal die aantal kapsules bepaal wat jou kind moet kry.

Jou dokter sal jou vertel hoe lank jou behandeling met CEZLIN sal duur. Moenie die behandeling vroegtydig stop nie. Indien jy die indruk het dat die effek van CEZLIN te sterk of te swak is, vertel jou dokter of apteker.

Neem CEZLIN saam met kos en drank
CEZLIN kan saam met kos en water geneem word.

As jy meer CEZLIN neem as wat jy moet
In die geval van oordosis, raadpleeg jou dokter of apteker sonder versuim. Indien nie een beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum.

As jy vergeet om CEZLIN te neem
Dit is belangrik dat jy voortgaan om CEZLIN te neem volgens die instruksies op die etiket en dat jy geen dosisse mis nie. As jy egter 'n dosis mis, neem dit so gou as wat jy onthou en neem die volgende dosis op die normale tyd. Moenie 'n dubbeldosis neem om vergete individuele dosisse in te haal nie.

Effekte as CEZLIN gestop word
Jy moenie ophou om CEZLIN te neem nie, tensy jou dokter jou beveel om dit te doen. As jy enige verdere vrae het oor die gebruik van CEZLIN, vra jou dokter of apteker.

4. Moontlike newe-effekte
CEZLIN kan newe-effekte hê.
Nie alle newe-effekte wat vir CEZLIN aangemeld is, is in hierdie pamflet ingesluit nie. Indien jou algemene gesondheid versleg of as jy enige nadelige effekte ervaar terwyl jy CEZLIN neem, raadpleeg asseblief jou dokter, apteker of ander gesondheidswerker vir raad.

Indien enige van die volgende gebeur, hou op om CEZLIN te neem en vertel jou dokter dadelik of gaan na die ongevalle-afdeling by jou naaste hospitaal:
- Inflammasie in die derm, los stoelgange en buikpyn, hipersensitiwiteit, SJS (ernstige siekte met uiserasie van die mond, lippe en vel), TEN (ernstige siekte en afskilfering van die vel), AGEP (velreaksies), DRESS ('n uitslag, betrokkenheid van interne organe, hematologiese abnormaliteite en sistemiese siektes)
- erge, aanhoudende of bloederige diarree (wat met maagpyn of koors geassosieer kan word). Dit is 'n algemene newe-effek wat kan voorkom na behandeling met antibiotika en kan 'n teken wees van ernstige dermontsteking (pseudomembraneuse-kolitis)
- tekens van 'n ernstige allergiese reaksie soos skielike hyg, moeilike asemhaling, swelling van ooglede, gesig of lippe, keel of tong, uitslag of jeuk (wat veral die hele liggaam aantas)
- 'n wydverspreide uitslag met blase en afskilfering van groot dele van die vel, veral rondom die mond, neus, oë of geslagsdele, bekend as Stevens-Johnson-sindroom, of toksiese epidermale nekrolise
- 'n seldsame veluitbarsting wat gekenmerk word deur die vinnige verskyning van areas van rooi vel besaai met klein pustules (klein blase gevul met wit/geel vloeistof) (kute veralgemeende eksantematiese pustulose (AGEP - Acute Generalised Exanthematous Pustulosis))
- veluitslag, wat blase kan vorm, en lyk soos klein teikens (sentrale donker kolle omring deur 'n ligter area, met 'n donker ring om die rand - eritherna multiforme)
- wydverspreide rooi veluitslag met klein etterbevattende blase (bulleuse eksfoliatiewe dermatitis)
- koors, geswelde limfknope of veluitslag, dit kan simptome wees van 'n toestand bekend as DRESS (Dwelmreaksie met Eosinofilie en Sistemiese Simptome)

Dit is alles baie ernstige newe-effekte. As jy dit het, het jy dalk 'n ernstige reaksie teenoor CEZLIN gehad. Jy mag dalk dringende mediese aandag of hospitalisasie benodig.

Vertel jou dokter dadelik of gaan na die ongevalle-afdeling by jou naaste hospitaal as jy enige van die volgende opmerk:
- Vergeling van die vel en oë, donker urine en moegheid wat simptome van lewerprobleme kan wees.

Dit is alles ernstige newe-effekte. Jy benodig dalk dringende mediese hulp.

Vertel jou dokter as jy enige van die volgende opmerk:
Gereelde newe-effekte
- Braking
- Naarheid (siek voel)
- Buikkrampe of maagpyn
- Uitslag makulo-papulêr (teenwoordigheid van plat en papules uitslag)
- Urtikaria jeukerige uitslag of swellings

Minder gereelde newe-effekte
- *Clostridium difficile* kolitis (ligte diarree tot dodelike kolitis)
- vaginale infeksie
- Agranulositose (tekort aan granulosiete in die bloed)
- Neutropenie ('n lae vlak van neutrofiele)
- Eosinofilie ('n toename in die aantal eosinofiele in die bloed)
- Leukopenie ('n vermindering in die aantal leukosiete) bloedselle wat veg vir infeksie in die bloed
- Trombositopenie ('n vermindering in die aantal bloedplaatjies in die bloed)
- 'Dysgeusia' (smaakversteuring)
- Esofageale ulkus
- Esofagitis (ontsteking van die omhulsel van die slukderm)
- Pruritus (jeukerige vel)
- morbilliforme uitslag (veluitslag)

Indien jy enige newe-effekte opmerk wat nie in hierdie pamflet genoem word nie, stel asseblief jou dokter of apteker in kennis.

Rapportering van newe-effekte
As jy newe-effekte kry, praat met jou dokter, apteker of verpleegster. Dit sluit enige moontlike newe-effekte in wat nie in hierdie pamflet gelys word nie.
Jy kan ook newe-effekte via https://www.sahpra.org.za/Publications/Index/8 aanmeld.
Deur newe-effekte aan te meld, kan jy help om meer inligting oor die veiligheid van hierdie medisyne te verskaf.

5. Hoe om CEZLIN te berg
- BERG ALLE MEDISYNE BUITE BEREIK VAN KINDERS.
- Berg teen of onder 25 °C.
- Hou in die oorspronklike verpakking tot benodig vir gebruik.
- Moenie na die vervaldatum, soos op die etiket, gebruik nie.
- Neem alle ongebruikte medisyne terug na jou apteker.
- Moenie ongebruikte medisyne in dreine of rioolstelsels (bv. toilette) weggooi nie.

6. Inhoud van verpakking en ander inligting
Wat CEZLIN bevat
Die aktiewe bestanddeel in CEZLIN is klindamisienhidrochloried USP gelykstaande aan klindamisien 150 mg.

Die ander bestanddele is
Laktosemonohidraat
Meelestysel
Talk
Magnesiumstearaat
Leë harde gelatienkapsule-dop (ligblou ondeursigtig / liggroen deursigtige grootte '1')

Hoe lyk CEZLIN en die inhoud van die verpakking
Ligblou ondeursigtig / Liggroen deursigtige grootte '1' harde gelatienkapsule gevul met wit tot naaswit poeier.

HDPE-houerpak (20's)
Wit ondeursigtige ronde 40 ml HDPE-houer met 33 mm nekafwerking toegemaak met 33 mm wit ondeursigtige polipropileen RS-sluiting met induksie-seëlvoering. Elke HDPE-houer bevat ook 1 g silikagel-sakkie. Die HDPE-bottels is verder verpak in voorafgedrukte kartonhouers met 'n pamflet.

HDPE-houerpak (100's)
Wit ondeursigtige ronde 120 ml HDPE-houer met 38 mm nekafwerking toegemaak met 38 mm wit ondeursigtige polipropileen CR-sluiting met induksie-seëlvoering. Elke HDPE-houer bevat ook 1 g silikagel-sakkie. Die HDPE-bottels is verder verpak in voorafgedrukte kartonhouers met 'n pamflet.

Naam, besigheidsadres en telefoonnommer van die houer van die sertifikaat van registrasie
Novagen Pharma (Edms) Bpk
Kantoor 2, 100 Sovereignrylaan,
Route 21 Corporate Park,
Nellmapiusrylaan, Irene, 0157, Pretoria, Suid-Afrika
Tel. Nr.: +27 012 345 3175
Epos adres: info@novagenpharma.co.za

Registrasienommer
52/20.1.1/0028

Hierdie pamflet is laas hersien in
12 Oktober 2021

Die professionele inligting vir CEZLIN (150 mg) kapsules is beskikbaar op die Novagen Pharma webtuiste: http://www.novagenpharma.co.za/products/antibiotic/

P1538046

Bemark deur:
United Pharma Marketing (Edms) Bpk

2408A158