SCHEDULING STATUS

S1

PROPRIETARY NAME, STRENGTH AND PHARMACEUTICAL FORM

ADCO-NAPHENSYL 1 % (NOSE DROPS)

ADCO-NAPHENSYL 0,25 % (PAEDIATRIC NOSE DROPS)

Read all of this leaflet carefully because it contains important information for you. ADCO-NAPHENSYL is available without a doctor's prescription, for you to treat a mild illness. Nevertheless you still need to use ADCO-NAPHENSYL carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share **ADCO-NAPHENSYL** with any other person.
- Ask your pharmacist if you need more information or advice.
- You must see your doctor if your symptoms worsen or do not improve after 10 days.

WHAT ADCO-NAPHENSYL CONTAINS

The active substance is phenylephrine hydrochloride.

Each 10 ml **ADCO-NAPHENSYL** 1 % contains 100 mg phenylephrine hydrochloride

Inactive ingredients: sodium chloride, sodium citrate and propylene glycol .

Contains benzalkonium chloride (0,022 % m/v) and sodium metabisulphite (0,1 % m/v) as preservatives.

Each 10 ml **ADCO-NAPHENSYL 0,25 %** contains 25 mg phenylephrine hydrochloride.

Inactive ingredients: sodium chloride, sodium citrate and propylene glycol .

Contains benzalkonium chloride (0,022 % m/v) and sodium metabisulphite (0,1 % m/v) as preservatives.

WHAT ADCO-NAPHENSYL IS USED FOR

ADCO-NAPHENSYL is used for temporary stuffiness of the nose caused by hay fever, allergies, or sinus episodes.

BEFORE YOU USE ADCO-NAPHENSYL

You should not use ADCO-NAPHENSYL:

- **if you are allergic (hypersensitive) to phenylephrine or to any of the ingredients of ADCO-NAPHENSYL.**
- **if your child is younger than 2 years.**
- **If you are** hypersensitive (allergic) or have overactive thyroid, heart or blood vessel disease, high blood pressure.
- if you are using monoamine oxidase inhibitor type antidepressant medication such as tranyclcypromine sulphate or moclobemide or within 14 days of stopping such treatment.
- if you are using tricyclic type antidepressants such as clomipramine, imipramine, amitryptiline or dothiepin.
- if you are taking rasagiline or selegiline.
- if you are taking reserpine and methyldopa for high blood pressure
- Ask your doctor or pharmacist if you are unsure of any of the above applies to you.

Take special care before using ADCO-NAPHENSYL:

Frequent or prolonged application will cause re-occurrence of the original symptoms by a rebound action of the lining of the nose in some cases. Overusage can also cause local irritation or allergy.

Pregnancy and breastfeeding:

If you are pregnant or breastfeeding your baby, please consult your doctor, pharmacist or other healthcare professional before using this medicine.

Using other medicines:

Always tell your healthcare professional if you are taking any other medicine. (This includes complementary or traditional medicines.)

Certain medicine can influence the mechanism of action of ADCO-NAPHENSYL and vice versa.

HOW TO USE ADCO-NAPHENSYL

Do not share medicines prescribed for you with any other person.

The usual dose is to instil two or three drops into the nose every four hours.

If you use more ADCO-NAPHENSYL than you should:

In the event of overdosage, consult your doctor, pharmacist or professional healthcare provider. If neither is available seek help at the nearest hospital or poison control centre.

The symptoms reported for overdosage in adults and children: are headache, vomiting, sweating, shaking or fainting.

POSSIBLE SIDE EFFECTS

Not all the side effects reported for ADCO-NAPHENSYL are included in this leaflet. Should your general health worsen or if you experience untoward effects while using ADCO-NAPHENSYL please consult your doctor, pharmacist or healthcare professional for advice.

The following side effects may occur less frequently: High blood pressure, vomiting, headache local irritation, allergic reaction.

If you notice any side effects not mentioned in the leaflet, please inform your doctor or pharmacist.

STORING AND DISPOSING OFADCO-NAPHENSYL

KEEP ALL MEDICINE OUT OF THE REACH OF CHILDREN.

Store at or below 25 °C. Protect from light and keep tightly closed.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains, sewerage systems (e.g. toilets).

PRESENTATION OF ADCO-NAPHENSYL

Dropper bottle containing 10 ml.

IDENTIFICATION OF ADCO-NAPHENSYL

ADCO-NAPHENSYL 1 %: A clear, colourless liquid.

ADCO-NAPHENSYL 0,25 %: A clear, colourless liquid.

REGISTRATION NUMBERS

ADCO-NAPHENSYL 1 %: H1403 (Act 101/1965)

ADCO-NAPHENSYL 0,25 %: H1596 (Act 101/1965)

NAME AND ADDRESS OF THE HOLDER OF THE REGISTRATION CERTIFICATE

Adcock Ingram Limited

1 New Road

Erand Gardens

Midrand, 1685

Private Bag X69

Bryanston, 2021

www.adcock.com

DATE OF PUBLICATION

Date of registration:

H1596 (Act 101 of 1965)

H1403 (Act 101 of 1965)

Date of revision: 17 February 2017