SILICONE GEL SHEET
(For Scar Management)

INSTRUCTIONS FOR USE

PRODUCT DESCRIPTION:

NOMOSCAR is a soft, clear, transparent, comfortable, re-usable self-adhesive sheet made of medical grade silicone, adhered on release liner. **NOMOSCAR** is supplied non-sterile in medical grade peel open pouches. The product can be cut to size depending on the size of the scar. **NOMOSCAR** is biocompatible as per ISO 10993 standards.

INDICATIONS/INTENDED USE:

NOMOSCAR is indicated for use in the prevention and/or in the management of old and newly formed, red and raised scars (hypertrophic scars/keloids).

METHOD OF APPLICATION:

1. Wash hands before use.
2. Clean the scar and surrounding skin with a mild soap and thoroughly dry the scar and surrounding skin.
3. Peel open the pouch and remove the **NOMOSCAR** from the pack.
4. For small scars, cut **NOMOSCAR** with clean scissors to fit the scar size, with a minimum of 1 cm overlap over the surrounding intact skin.
5. For large scars, use as many **NOMOSCAR** sheets next to each other as required to cover the entire scar, with a minimum of 1 cm overlap over the surrounding intact skin.
6. Store the remaining portion of the unused **NOMOSCAR** in the product pouch in a dry place.
7. Remove the release liner with care to expose the adhesive side of the **NOMOSCAR**.
8. Apply **NOMOSCAR** adhesive side to the scar without stretching.
9. For the first two days, use **NOMOSCAR** for up to 4 hours, and up to 8 hours for the next two days. The usage can be increased incrementally by 2 hours per day until a maximum of 18 hours per day is reached.
10. Wash **NOMOSCAR** after every use. Apply mild soap on the adhesive side of **NOMOSCAR** and keep under gently flowing tap water. Ensure to remove all the dirt (dust, hair, fabric strings, etc.) and dry with a lint free cloth.
11. Dry your hands thoroughly and place the **NOMOSCAR** on the release liner.
12. During re-application, if necessary, an adhesive tape can be used to keep **NOMOSCAR** in place.
13. During application of **NOMOSCAR**, if the adhesive side sticks to itself, just hold it under flowing tap water and carefully detangle it.
14. When **NOMOSCAR** begins to chip off or has lost its adhesive property completely or if removal of dirt is difficult, it should be replaced with a new piece of **NOMOSCAR**. This usually occurs after 22 to 25 times of application.

CONTRAINDICATIONS:

1. Do not use on open wounds.
2. Do not use on skin until suture removal is complete.
3. Do not use on fragile skin affected by dermatological conditions (e.g. psoriasis, eczema, etc.).
4. Do not use any ointments or creams under **NOMOSCAR**.
5. Do not use **NOMOSCAR** on patients with known hypersensitivity to silicone.

PRECAUTIONS & WARNINGS:

1. If rash appears, reduce **NOMOSCAR** treatment time per day and observe. If rash persists, discontinue treatment and consult your doctor.
2. **NOMOSCAR** is strictly for external use only.
3. Keep out of reach of children.

PRODUCT AVAILABILITY:

NOMOSCAR is supplied non-sterile and is available in medical grade peel open pouch, packed in the following dimensions:
3 cm x 15 cm.
10 cm x 10 cm.

STORAGE:

Store in a dry place.

SHELF LIFE:

NOMOSCAR has a shelf life of three years from the date of manufacturing.

DEVICE DISPOSAL:

Disposal should be handled according to local environmental procedure.

LP0040/0002

Marketed in South Africa by:

MediVice
Medical Device Division of
LeBasi Pharmaceuticals (Pty) Ltd
087 551 3245 I www.lebasi.co.za

San Domenico Building,
Unit 6, 10 Church Street,
Durbanville, Cape Town 7551

Under licence from:

CORTIS MEDTECH PTE. LTD.
#05-03 plus, 20 Cecil Street, Singapore 049705