1.3.2 PATIENT INFORMATION LEAFLET

SCHEDULING STATUS

S2

PROPRIETARY NAME AND DOSAGE FORM

PANAMOR 50 mg DISPERSIBLE TABLET

Read all of this leaflet carefully because it contains important information for you

PANAMOR 50 mg DISPERSIBLE TABLET

is available without a doctor's prescription, for you to treat a mild illness.

Nevertheless, you still need to use PANAMOR 50 mg DISPERSIBLE TABLET carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share PANAMOR 50 mg DISPERSIBLE TABLET with any other person.
- Ask your pharmacist if you need more information or advice.
- You must see a doctor if your symptoms worsen or do not improve after 3 days.

WHAT PANAMOR 50 mg DISPERSIBLE TABLET CONTAINS

Each PANAMOR 50 mg DISPERSIBLE TABLET contains 46,5 mg of diclofenac free acid.

Other ingredients are:

Croscarmellose sodium, glyceryl dipalmitostearate, microcrystalline cellulose, purified talc, sodium saccharin, sodium starch glycollate.

Contains sweetener: Sodium saccharin 0,26 mg

WHAT PANAMOR 50 mg DISPERSIBLE TABLET IS USED FOR

This medicine is used to treat gout attacks.

BEFORE YOU TAKE PANAMOR 50 mg DISPERSIBLE TABLET

In deciding to use a medicine, the risk of taking the medicine must be weighed against the good it will do. Consider the following before deciding to take this medicine:

- Tell your healthcare professional if you have had any allergic reaction to any medicines, food or even a preservative.
- Tell your healthcare professional if you are pregnant or breastfeeding. This product should not be used in pregnancy or breastfeeding mothers.
- Certain medicines may not be used together. It is especially important that you tell your healthcare professional if you are taking any of the following: anticoagulants, methotrexate, lithium, potassium supplements, digoxin or anti-diabetic therapy.

- Other medical conditions may affect the use of this medicine. Inform your healthcare professional if you have any other medical problem, especially liver or kidney disease, stomach ulcer or other intestinal problem, diabetes mellitus, anaemia, bleeding problems

HOW TO TAKE PANAMOR 50 mg DISPERSIBLE TABLET

Take this medicine exactly as your doctor, pharmacist or healthcare professional told you.

Dissolve one tablet in water and take every 8 hours (three times a day) for acute gout attacks subject to a maximum daily dose of 150 mg for a maximum treatment period of 3 days (three PANAMOR 50 mg DISPERSIBLE TABLET in divided doses).
Do not take more than three tablets a day.

Use the lowest effective dose for the shortest possible duration of treatment.

These tablets cannot be swallowed whole. Place a tablet in a glass full of water, the tablet will disperse, once dispersed fully, drink. If there is medicine on the inside of the glass rinse the glass with water and drink.

If you have the impression that the effect of PANAMOR 50 mg DISPERSIBLE TABLET is too strong or too weak, tell your doctor or pharmacist.

SIDE EFFECTS

PANAMOR 50 mg DISPERSIBLE TABLET can have side effects.

Not all side effects reported for PANAMOR 50 mg DISPERSIBLE TABLET are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking PANAMOR 50 mg DISPERSIBLE TABLET, please consult your healthcare provider for advice.

Along with its needed effects, a medicine may cause some unwanted effects. Some common side effects that may not need medical attention include, abdominal cramps, diarrhoea, indigestion, headache, nausea or vomiting. Insomnia, depression, or ringing sound in the ears should be reported. Seek medical attention if any of the following effects occurs tightness of chest, irregular heartbeat, hive-like swellings on face, eyelids, mouth, lips or tongue.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

STORING AND DISPOSING OF PANAMOR 50 mg DISPERSIBLE TABLET

Store at or below 25°C in tightly closed containers.

Protect from light and moisture.

Keep in original packaging until required for use.

Do not store in a bathroom.

Do not use after expiry date stated on the label.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

STORE ALL MEDICINES OUT OF REACH OF CHILDREN.

PRESENTATION OF PANAMOR 50 mg DISPERSIBLE TABLET

9 tablets are packed in a clear polyvinyl chloride and polyethylene rigid film with a transparent polyvinylidene copolymer coating sealed with an aluminium backing foil. One strip of 9 tablets is packed into an outer cardboard carton with a leaflet.

IDENTIFICATION OF PANAMOR 50 mg DISPERSIBLE TABLET

A white, diamond-shaped biconvex tablet bisected on one side and debossed with “P” on the other side.

REGISTRATION NUMBER

A38/3.1/0383

NAME, BUSINESS ADDRESS AND TELEPHONE NUMBER OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION

PHARMACARE LIMITED

Healthcare Park

Woodlands Drive

Woodmead 2191

Hotline: 0800 122 912 (South Africa)

Tel: +27 11 239 6200 (Other)

DATE OF PUBLICATION

Date of registration: 23 February 2005

The date of the most recent amendment to the patient information leaflet as approved by the Authority: 23 February 2005

ZA_PANADTAB_0502_01