WHAT TRANSACT CONTAINS:

TransAct patches are white polyester based with a menthol-type aroma and contains 40 mg Flurbiprofen.

Other ingredients include carmellose sodium, concentrated Glycerin, crotamiton, dried aluminium hydroxide gel, emulsion of methyl acrylate and 2-ethylhexyl acrylate copolymer, i-menthol, isopropyl myristate, mentha oil *(peppermint Oil)*, partially neutralised polyacrylate, titanium dioxide E171 *(CI no. 77891)*, polysorbate 80, polypropylene film, water, sorbitan sesquioleate, starch grafted acrylate 300, tartaric acid, talc, unwoven polyester fabric.

WHAT THIS MEDICINE IS USED FOR:

TransAct is indicated for the symptomatic relief of localised pain and inflammation associated with sprains, strains and muscular or arthritic conditions.

BEFORE YOU APPLY TRANSACT:

You must consult your doctor or pharmacist before using TransAct if:

- You are pregnant, planning to become pregnant or breast feeding.
- You suffer from liver, kidney or heart disease.
- You suffer from asthma or have an allergic reaction or wheezing after taking aspirin or other arthritis medicine.
- You are taking any of the following medicines – aspirin or other drugs used to treat arthritis.
- You are on medicines for thrombosis *(eg. Warfarin)*, blood pressure or water tablets.
- You have previously shown allergic reaction to the active ingredient, Flurbiprofen.
- You have a history of non-allergic asthma.
- You have a history of stomach or duodenal ulcers, internal bleeding of stomach or intestines, a chronic disease affecting the colon *(large bowel)*, heart failure or high blood pressure.

If the answer is yes to any of these questions, tell your doctor or pharmacist.

DO NOT USE TRANSACT:

- On broken skin, fragile or skin affected by irritation or infection

Systemic absorption *(dosage of the product absorbed)* is increased if this product is applied to damaged skin.

- In children younger than 12 years of age.
- Continuously for more than 4 weeks.

Always tell your doctor, pharmacist or healthcare provider if you are taking any other medicines on a regular basis, concomitant use of these medicines may cause undesirable interactions.

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby please consult your doctor, pharmacist or other health care provider for advice before using this medicine

HOW TO APPLY TRANSACT PATCHES:

Do not share medicines prescribed for you with any other person.

Always use TransAct exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.


TransAct is for external use only. Use only one patch at a time on your skin and replace every 12 hours.



Wipe the affected area clean.



Remove one patch from the sachet and ensure that the sachet is securely re-closed.



First rub the corner of the patch between the thumb and fingers to peel off the backing and apply the adhesive side to the skin.



When applying stretch the patch gently to prevent the surface of the patch wrinkling. When first applied this product may feel cool.



Where a joint is affected slightly bend the joint before applying.



You may find it helpful to use a bandage to keep the patch in place on joints like a knee or elbow.

It is recommended that bathing or showering should be arranged to coincide with routine changing of the patch.

Do not get the patch wet – remove before bathing.

©2020. TransAct is a registered trademark.

POSSIBLE SIDE EFFECTS:

Not all side-effects reported for this medicine are included in this leaflet.

Should your general health worsen or if you experience any untoward effects while taking this medicine, please consult your doctor, pharmacist or other health care professional for advice.

This medicine sometimes causes side effects in some people such as local itching, redness, numbness and tingling. If these becomes troublesome, consult your doctor. If you get any of the following, stop using the medicine and tell your doctor: Nausea, diarrhoea, rashes or gastric upset.

When TransAct is used for long periods of time, the possibility of side effects throughout the body cannot be completely excluded. These side-effects which you may experience with Flurbiprofen *(the active ingredient in your medicine)* include the following: Pain in the upper, middle region of the stomach, belching, nausea *(queasiness)* and diarrhoea *(frequent or loose watery stools)* headache or slight dizziness. If any of the above-mentioned side-effects continue or are troublesome, you must stop using the medicine and consult your doctor or pharmacist.

There have also been reports of skin rash, swelling of the hands and feet due to retained fluid, the formation of an ulcer of the stomach or intestines, internal bleeding of the stomach or intestines, allergic reactions *(e.g. constriction of the air passages and severe allergic reaction)*, yellow pigmentation of the skin and eyes, disease of the liver or kidney and kidney failure.

Isolated cases of the erythema multiforme *(skin condition)* have been observed and present in the following manner: skin lesions, itching, joint pain and fever. During prolonged treatment with TransAct, blood counts and monitoring of kidney and liver function are indicated as precautionary measures.

STORING AND DISPOSING OF TRANSACT:



STORE ALL MEDICINES OUT OF THE REACH OF CHILDREN.

Store at or below 25 °C and reseal the sachet after use.

Do not use this product after the expiry date shown on the carton and the back of each sachet.

Do not use any patches remaining one month after opening the sachet.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems *(e.g. toilets)*.

PRESENTATION:

Carton containing either 1 or 2 resealable laminated sachets.

Each sachet contains 5 TransAct patches.

IDENTIFICATION:

A 10 X 14 cm patch consisting of an un-woven polyester backing spread evenly with a white to pale yellow ointment and covered by a peel-off liner.

The ointment has an odour of peppermint.

REGISTRATION NUMBER:

28/3.1/0268

NAME, ADDRESS AND TELEPHONE NUMBER OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:

Reckitt Benckiser Pharmaceuticals (Pty) Ltd.
8 Jet Park Road
Elandsfontein 1601
+2711 871 1611

DATE OF PUBLICATION OF THE PATIENT INFORMATION LEAFLET:

Date on the registration certificate: 27 May 1994
Date of the revised patient information leaflet: 26 May 2020

TransAct®

TransAct®

Medisinale Pynverligtings Plakker
Vir Spier- en Gewrigspyn

40 mg Flurbiprofeen
per plakker

SKEDULERINGSTATUS:

S0

EIENDOMSNAAM, STERKTE EN FARMASEUTIESE VORM:

TransAct®: Flurbiprofen 40 mg per pleister.
Transdermale Terapeutiese Stelsel.

Lees die hele pamflet sorgvuldig, want dit bevat belangrike inligting.

TransAct is sonder 'n voorskrif van 'n dokter beskikbaar om ligte siekte te behandel. U moet nietemin steeds TransAct versigtig gebruik om die beste resultate te verkry.

Bewaar hierdie pamflet, ingeval u dit weer moet lees.
Moenie TransAct met enige ander persoon deel nie.
Vra u apteker as u meer inligting of advies benodig.

Professionele inligtingspamflet is beskikbaar by https://www.cpdcentre.co.za

WAT TRANSACT BEVAT:

TransAct pleister is gemaak van wit polyester, met ‘n menthol-aroma en bevat 40 mg Flurbiprofen.

Ander bestanddele sluit in: karmellose-natrium, gekonsentreerde gliserien, crotamiton, gedroogde aluminiumhidroksiedgel, emulsie van metielakriellaat en 2-etielheksielakriellaat-copolymeer, i-menthol, isopropylmyristaat, mentha-olie *(pepermentolie)*, gedeeltelik geneutraliseerde poli-akriellaat, titaandioksied E171 (*C no. 77891)*, polysorbaat 80, polipropileen film, water, sorbitan sesquioleate, stysel-ent akriellaat 300, wynsteensuur, talk, ongeweefde polyester materiaal.

WAARVOOR HIERDIE MEDISYNE GEBRUIK WORD:

TransAct word aangedui vir die simptomatiese verligting van gelokaliseerde pyn en ontsteking wat verband hou met verstuitings, spanning en spier- of artritis-toestande.

VOORDAT U TRANSACT AANWEND:

U moet u dokter of apteker raadpleeg voordat u TransAct gebruik indien:

- U swanger is, beplan om swanger te raak of borsvoed.
- U aan lewer-, nier- of hartsiektes ly.
- U aan asma ly, of ‘n allergiese reaksie kry of kortasem raak nadat u aspirien of ander artritismedisyne geneem het.
- U een van die volgende medisynes neem - aspirien of ander medisyne wat gebruik word om artritis te behandel.
- U medisyne vir trombose *(bv. Warfarin)*, bloeddruk of watertablette neem.
- U voorheen ‘n allergiese reaksie op die aktiewe bestanddeel, Flurbiprofen, getoon het.
- U ‘n geskiedenis van nie-allergiese asma het.
- U ‘n geskiedenis van maagswere of duodenale swere, interne bloeding van die maag of derms, enige chroniese siekte wat die dikderm *(kolon)* affekteer, hartversaking of hoë bloeddruk het.

As die antwoord op enige van hierdie vrae ja is, laat weet u dokter of apteker.

MOENIE TRANSACT IN ENIGE VAN DIE VOLGENDE GEVALLE GEBRUIK NIE:

- Op vel wat gebreek, broos, of deur irritasie of infeksie geaffekteer word.

As die produk op beskadigde vel toegedien word, word sistemiese opname *(dosis van die geabsorbeerde produk)* verhoog.

- By kinders jonger as 12 jaar.
- Deurlopend vir meer as 4 weke.

Laat weet altyd u dokter, apteker of gesondheidsorgverskaffer indien u gereeld ander medisyne neem, aangesien die gelyktydige gebruik van hierdie medisyne ongewenste interaksies kan veroorsaak.

As u swanger is of borsvoed, dink dat u swanger mag wees of van plan is om ‘n baba te hê, raadpleeg u dokter, apteker of ander gesondheidsorgverskaffer voordat u hierdie medisyne gebruik.

HOE OM TRANSACT PLEISTERS AAN TE WEND:

Moet nooit medisyne wat vir u voorgeskryf is, met enige ander persoon deel nie.

Gebruik altyd TransAct presies soos in hierdie pamflet beskryf word, of soos u dokter, apteker of verpleegkundige aanbeveel. Raadpleeg u dokter, apteker of verpleegkundige as u onseker is.


TransAct is slegs vir eksterne gebruik. Gebruik slegs een pleister op ‘n slag op die vel en vervang elke 12 uur.



Vee die geaffekteerde area skoon.



Haal een pleister uit die pakkie en maak seker dat die pakkie weer goed geseël is.



Vryf eers die hoek van die pleister tussen die duim en vingers om die rugkant los te maak, en wend die kleefkant op die vel aan.



Wanneer die pleister aangeend word, rek dit liggies om te voorkom dat die oppervlak van die pleister verrimpel. Wanneer die produk aanvanklik aangewend word, kan dit koel voel.



Waar ‘n gewrig geaffekteer is, buig die gewrig effens voordat die pleister aangewend word.



U mag dit nuttig vind om ‘n verband te gebruik om die pleister op gewrigte, soos ‘n knie of elmboog, in plek te hou.

Dit word aanbeveel dat u u bad of storttyd reël rondom die tyd wat die pleister vervang moet word.

Moenie die pleister laat nat word nie - verwyder voordat u bad.

©2020. TransAct is ‘n geregistreerde handelsmerk.

MOONTLIKE NEWE-EFFEKTE:

Nie alle newe-effekte wat vir hierdie medisyne aangemeld word, is in hierdie pamflet opgeneem nie. Raadpleeg u dokter, apteker of ander gesondheidsorgwerker vir raad indien u algemene gesondheid vererger, of indien hierdie medisyne u nadelig beïnvloed.

Hierdie medisyne veroorsaak soms newe-effekte by sommige mense, soos plaaslike jeukerigeheid, rooiheid, gevoelloosheid en tinteling. As dit lastig raak, raadpleeg u dokter. As u een van die volgende probleme ondervind, moet u ophou om die medisyne te gebruik en u dokter laat weet: naarheid, diarree, uitslag of maagaandoening.

Wanneer TransAct vir lang periodes gebruik word, kan die moontlikheid van newe-effekte in die liggaam nie heeltemal uitgesluit word nie. Die newe-effekte wat u met Flurbiprofen *(die aktiewe bestanddeel in u medisyne)* kan ervaar , sluit die volgende in: Pyn in die boonste, middelste deel van die maag, winderigheid, naarheid en diarree *(gereelde of los, waterige ontlasting)*, hoofpyn of ‘n ligte duiseligheid. As een van die bogenoemde newe-effekte voortduur of lastig is, moet u ophou om die medisyne te gebruik en u dokter of apteker raadpleeg.

Daar was ook al berigte van veluitslag, swelling van hande en voete as gevolg van behoue vloeistof, die vorming van swere in die maag of ingewande, interne bloeding van die maag of ingewande, allergiese reaksies *(bv. vernouing van die lugweë en ernstige allergiese reaksie)*, geel pigmentasie van die vel en oë, lewer- of nier-siekte of -versaking.

Geïsoleerde gevalle van die erythema multiforme *(veltoestand)* is op die volgende manier waargeneem: velletsels, jeukerigheid, gewrigspyn en koors. Tydens langer behandelinge met TransAct, word bloedtellings en monitering van die nier- en lewerfunksie aangedui as voorsorgmaatreëls.

BERGING EN VERWYDERING VAN TRANSACT:



BÊRE ALLE MEDISYNE BUITE DIE BEREIK VAN KINDERS.

Bêre by of onder 25 ° C en seël die pakkie na gebruik.

Moenie die produk gebruik na die vervaldatum wat op die houer en agterkant van elke pakkie verskyn nie.

Moenie enige pleisters wat meer as ‘n maand na die pakkie oopgemaak is oorbly, gebruik nie.

Handig alle ongebruikte medisyne in aan u apteker.

Moenie van ongebruike medisyne in dreine of rioolstelsels *(bv. toilette)* ontslae raak nie.

AANBIEDING:

Kartonhouer wat óf 1 óf 2 herseëlbare, gelamineerde pakkies bevat.

Elke pakkie bevat 5 TransAct-pleisters.

IDENTIFIKASIE:

‘n Pleister van 10 x 14 cm, wat bestaan uit ‘n ongeweefde polyester-agterkant wat eweredig met ‘n wit tot liggeel salf gesmeer is en met ‘n aftrek-laag bedek is.

Die salf het ‘n pepermentreuk.

REGISTRASIENOMMER:

28/3.1/0268

NAAM, ADRES EN TELEFOONNOMMER VAN DIE SERTIFIKAAT VAN REGISTRASIEHOUER:

Reckitt Benckiser Pharmaceuticals (Pty) Ltd.
8 Jet Park Weg
Elandsfontein 1601
+2711 871 1611

PUBLIKASIEDATUM VAN DIE PASIËNTINLIGTINGSPAMFLET:

Datum op die registrasiesertifikaat: 27 Mei 1994
Datum van die hersiene pasiëntinligtingspamflet: 26 Mei 2020

S0 TransAct®

Medicated Pain Relief Patch

For Muscle & Joint Pain

40 mg Flurbiprofen per patch

SCHEDULING STATUS:

S0

PROPRIETARY NAME, STRENGTH AND PHARMACEUTICAL FORM:

TransAct®: Flurbiprofen 40 mg per patch.
Transdermal Therapeutic System.

Read all of this leaflet carefully because it contains important information for you.

TransAct is available without a doctor’s prescription, for you to treat a mild illness. Nevertheless you still need to use TransAct carefully to get the best results from it.

Keep this leaflet. You may need to read it again.

Do not share TransAct with any other person.

Ask your pharmacist if you need more information or advice.

Botswana	BOT1803319 (S3)
Namibia	04/3.1/0639 (NS1)
Zambia	133/016 (P)

Professional Information leaflet is available on https://www.cpdcentre.co.za

S0 TransAct®