

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS

Schedule 2

STUGERON®

25 mg Cinnarizine per tablet

Contains sugar (lactose monohydrate and sucrose)

Each tablet contains 158,8 mg lactose monohydrate and 15 mg sucrose

Read all of this leaflet carefully before you start using STUGERON

- Keep this leaflet. You may need to read it again.
- Do not share STUGERON with other people
- Ask your health care provider or pharmacist if you need more information or advice.
- You must see a doctor if your symptoms worsen or do not improve

What is in this leaflet

1. What STUGERON is and what it is used for
2. What you need to know before you take STUGERON
3. How to use STUGERON
4. Possible side effects
5. How to store STUGERON
6. Contents of the pack and other information

1. What STUGERON is and what it is used for

STUGERON is used to treat the following:

- **Balance disorder** - STUGERON relieves the dizziness and spinning sensation ("vertigo") caused by problems of the inner ear. It also relieves the associated

symptoms like persistent ringing in the ears (“tinnitus”), rapid and uncontrollable movements of the eyes, nausea and vomiting;

- STUGERON also prevents motion sickness.

2. What you need to know before you take STUGERON

Always inform your doctor or pharmacist if you are using other medicines because some medicines should not be taken together.

Do not take STUGERON if:

- you are hypersensitive (allergic) to cinnarizine or any of the ingredients of STUGERON
- you suffer from Parkinson’s disease.

Other medicines and STUGERON:

Always tell your healthcare professional if you are taking any other medicine. (This includes complementary or traditional medicines).

Some of the interactions are listed below:

- Medicines for depression, sleeping pills, tranquillisers and strong painkillers can have an increased effect when taken with STUGERON.
- Alcohol and STUGERON strengthen each other's sedative effect. You should not drink while on STUGERON.

Taking STUGERON with food and drink:

You should take STUGERON after meals. The tablets are best taken with a little water.

Pregnancy and Breastfeeding:

If you are pregnant or breastfeeding your baby, please consult your doctor, pharmacist or healthcare professional for advice before taking STUGERON

Driving and using machines

STUGERON may lead to drowsiness and impaired concentration. Especially at the start of treatment STUGERON can cause drowsiness, which can make you less alert and reduce your driving ability. So, you should be careful about using machines or driving while on STUGERON.

STUGERON contains lactose

STUGERON contains "lactose" (a type of sugar). Patients with rare hereditary conditions of lactose/fructose or galactose intolerance should not take STUGERON.

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking STUGERON.

STUGERON contains sugars (lactose and sucrose), which may have an effect on the control of your blood sugar if you have diabetes mellitus.

3. How to use STUGERON

Do not share medicines prescribed for you with any other person.

Always take STUGERON as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure.

Your doctor will tell you how much STUGERON you should take. It is best to start with a small amount of STUGERON, after which the dose can gradually be increased until the desired result is obtained. Your doctor will explain exactly how you should proceed.

In most cases one of the following doses is prescribed:

Disorders of balance-Adults: 1 x 25 mg tablet three times daily.

Prevention of Motion sickness:

Adults: 25 mg may be taken 2 hours before the start of the journey and 12,5 mg (half a tablet) to 25 mg may be repeated every 8 hours during the journey when necessary.

Children 8 to 12 years: 12, 5 mg (half a tablet) three times daily when necessary.

This formulation is not suitable for children less than 8 years old.

You should take STUGERON a half hour before you leave for a trip, and for long trips you can take it every 6 hours after that.

Important

The maximum recommended dosage should not exceed 225 mg (9 tablets) daily in adults- if necessary the dosage may be divided over 2 or 3 intakes per day. As the effect of STUGERON tablets on vertigo is dose dependent, the dosage should be increased progressively.

If you take more STUGERON than you should:

Taking too much STUGERON, can lead to the following signs and symptoms: alterations in consciousness ranging from sleepiness to loss of consciousness, vomiting, muscle weakness or incoordination, and convulsions. Death was reported in association with cinnarizine overdose.

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre.

4. Possible side effects

STUGERON can have side effects. Not all side effects reported for STUGERON are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking STUGERON, please consult your doctor, pharmacist or other healthcare professional for advice.

Listed below are side-effects (also called adverse drug reactions) related to treatment with STUGERON.

Frequent

- Sleepiness
- Weight gain
- Nausea

Less frequent

- Daytime sleepiness
- Vomiting, indigestion, stomach ache
- Excessive sweating
- Tiredness
- Liver problem that causes yellowing of the skin or eyes (jaundice)
- Itchy red lesions or rash on the skin or grey white pimples in the mouth
- Movement problems like jerky movements, muscle stiffness, trembling.
These symptoms are also known as “extrapyramidal symptoms” (EPS)
- Red circular lesions or rash that occur on sun exposed skin areas.
- Muscle stiffness

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist or nurse. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**” found online under SAHPRA’s publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of STUGERON.

5. How to store STUGERON

Store at or below 30°C (at normal room temperature). Protect from light.

Keep all medicines in their original packaging and in a dry place (never in the bathroom, for example).

STUGERON can be kept for only a limited period. Do not use STUGERON after the date (month and year) printed after "EXP", even if it has been stored properly.

Store all medicines out of reach of children.

How to Dispose of STUGERON

Return all unused medicine to your pharmacist. Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What STUGERON contains

The active substance is cinnarizine. Each tablet contains 25 mg of cinnarizine.

The tablet also contains: cotton seed oil hydrogenated, lactose monohydrate, maize starch, polyvidone, sucrose, talc.

Contains sugar (lactose monohydrate and sucrose)

Each tablet contains 158,8 mg lactose monohydrate and 15 mg sucrose.

What STUGERON looks like and contents of the pack

STUGERON is a white, circular, slightly arched, biconvex half-scored tablet, embossed with "S/25" on one side and "JANSSEN" on the other side.

Cartons contain one or more blister packs of 10, 20 or 25 tablets each.

Holder of Certificate of Registration



JANSSEN PHARMACEUTICA (PTY) LTD

(Reg. No. 1980/011122/07)

2 Medical Road, Halfway House

Midrand, 1685

Tel: +27 (0)11 518 7000

RA-JACZA-MedInfo@its.jnj.com

This leaflet was last revised in

- Date of registration: 13 May 1996
- Date of most recently revised Patient Information Leaflet as approved by Advisory Clinical Committee: 14 June 2022.

Registration number

C/5.7/590