PATIENT INFORMATION LEAFLET

Scheduling Status: S4

Proprietary Name, Strength and Pharmaceutical Form

MOMATE 0,1 % CREAM (each gram of cream contains 1 mg of Mometasone Furoate in a cream base)

- **Read all of this leaflet carefully before you start using MOMATE 0,1 % CREAM**
- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor or your pharmacist.

MOMATE 0,1 % CREAM has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

1. WHAT MOMATE 0,1 % CREAM CONTAINS

The active substance is mometasone furoate,

The other ingredients of the product are aluminium starch octenylsuccinate, hexylene glycol, phosphoric acid, phospholipon 90 H, purified water, titanium dioxide, white petrolatum and white wax.

2. WHAT MOMATE 0,1 % CREAM IS USED FOR

Mometasone belongs to a class of medicines known as corticosteroids.

MOMATE 0,1 % CREAM is used for the treatment of inflammation caused by psoriasis and is also used in skin conditions that respond to corticosteroids.

3. BEFORE YOU USE MOMATE 0,1 % CREAM

Do not use MOMATE 0,1 % CREAM:

- if you are allergic (hypersensitive) to mometasone furoate or any of the other ingredients of MOMATE 0,1 % CREAM
- if you have viral infections, such as herpes simplex, vaccinia and varicella or untreated fungal infections.
- if you are pregnant or breastfeeding.

Take special care with MOMATE 0,1 % CREAM:

MOMATE 0,1 % CREAM is not for use in the eyes.

If you accidentally get MOMATE 0,1 % CREAM in your eyes, flush it with water.

Use of MOMATE 0,1 % CREAM for periods longer than three weeks in children is not recommended.

Use of MOMATE 0,1 % CREAM in children under the age of two years is not recommended.

In facial dermatoses MOMATE 0,1 % CREAM should be used cautiously and for short period only. Do not bandage or otherwise wrap the skin being treated unless directed to do so by your doctor.

Please refer to your doctor regularly in cases where the product is used in large area for longer period, especially when it is used in children.

Do not use MOMATE 0,1 % CREAM in any skin crease area like face, groin or armpits unless directed to do so by doctor, and especially not in the nappy are of infants.

Do not use MOMATE 0,1 % CREAM for other skin problems without first checking with your doctor.

Do not use MOMATE 0,1 % CREAM more often or for longer time than your doctor instructed.

Stop using MOMATE 0,1 % CREAM and seek immediate medical attention if irritation or sensitisation occurs and treat appropriately.

Pregnancy and Breastfeeding:
MOMATE 0,1 % CREAM should not be used in pregnancy or breastfeeding.
If you are pregnant or breastfeeding your baby please consult your doctor, pharmacist or other healthcare professional for advice before using MOMATE 0,1 % CREAM

Driving and using machinery
MOMATE 0,1 % CREAM has no side effects such as dizziness or drowsiness, which may affect the ability to drive or operate machinery.

Taking / Using other medicines with MOMATE 0,1 % CREAM
Always tell your healthcare professional if you are taking/using any other medicine. (This includes complementary or traditional medicines.)

4. HOW TO USE MOMATE 0,1 % CREAM:
Do not share medicines prescribed for you with any other person.
Always apply MOMATE 0,1 % CREAM exactly as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure.
MOMATE 0,1 % CREAM is not recommended for use in children under two years.

If you forget to use MOMATE 0,1 % CREAM:
Try to apply your medicine at the same time every day. If you miss a dose, apply it as soon as possible unless it is nearly time to apply again.

Effects when treatment with MOMATE 0,1 % CREAM is stopped:
Do not stop applying MOMATE 0,1 % CREAM without first consulting your doctor.
If you have any further questions on the use of MOMATE 0,1 % CREAM ask your doctor or pharmacist.

5. POSSIBLE SIDE EFFECTS:

MOMATE 0,1 % CREAM can have side effects.

Long term use may cause changes in the skin, leading to: thinning of the skin, loss of elasticity, dilation of blood vessels, bruising, and red vein-like spots on the skin and around the mouth. These effects occur especially when obstructing bandages are used.

Less frequent:

Hormone related disorders – growth retardation, most likely severe in children.

Skin and tissue disorders - allergic eczema, burning, dryness, pustules around the hair strands, irritation, itching, secondary infection, thinning of the skin, acne-like sores, excessive hair growth, lightening of the skin, sweat rash, wrinkling of the skin and stripes.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

6. STORING AND DISPOSING OF MOMATE 0,1 % CREAM:

Store all medicines out of reach of children.

Store at or below 25 °C. Do not refrigerate.

Keep well closed and protect against light.

For external use only.

Do not use after the expiry date stated on the carton

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets)

7. PRESENTATION OF MOMATE 0,1 % CREAM:

MOMATE 0,1 % CREAM is packaged in 20 g laminated closed end nozzle aluminium tube with milky white cap having a piercing point

8. IDENTIFICATION OF MOMATE 0,1 % CREAM:

Smooth, white to off-white cream, free from grittiness.

9. REGISTRATION NUMBER

43/13.4.1/0250

10. NAME AND ADDRESS OF REGISTRATION HOLDER

Activo Health (Pty) Ltd

Block B, Arena Office Park

272 West Avenue

Centurion

0157

11. DATE OF PUBLICATION

29 May 2023