SCHEDULING STATUS: S0

PROPRIETARY NAME, STRENGTH AND DOSAGE FORM

STREPSILS STRAWBERRY LOZENGES, 2,4-dichlorobenzyl alcohol 1,2 mg and amylmetacresol 0,6 mg.

Read all of this leaflet carefully because it contains important information for you

Strepsils Strawberry Lozenges is available without a doctor's prescription, for you to treat a mild illness. Nevertheless, you still need to take Strepsils Strawberry Lozenges carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share Strepsils Strawberry Lozenges with any other person.
- Ask your pharmacist if you need more information or advice.
- You must see a doctor if your symptoms worsen or persist or do not improve after 2 days

1. WHAT STREPSILS STRAWBERRY LOZENGES CONTAIN

- The active substances are 2,4-dichlorobenzyl alcohol and amylmetacresol. Each lozenge contains 2,4-dichlorobenzyl alcohol 1,2 mg and amylmetacresol 0,6 mg.
- The other ingredients are: Flav P Strawberry flavour 052312B, , liquid, pink antho P-WS (E163), purified water and tartaric acid.
- Contains sweeteners (isomalt 1830 mg, maltitol 481 mg, sodium saccharin 2 mg per lozenge)

2. WHAT STREPSILS STRAWBERRY LOZENGES ARE USED FOR

- Strepsils Strawberry Lozenges contain antiseptic ingredients that are used for the relief of minor throat and mouth infections as well as sore throats

3. BEFORE YOU TAKE STREPSILS STRAWBERRY LOZENGES

Do not take Strepsils Strawberry lozenges:

If you are hypersensitive (allergic) to 2,4-dichlorobenzyl alcohol, amylmetacresol, or to any of the other ingredients of Strepsils Strawberry Lozenges

Take special care with STREPSILS STRAWBERRY LOZENGES

STREPSILS STRAWBERRY LOZENGES contains isomaltitol and maltitol syrup, which may have a mild laxative effect if several are taken a day. Patients with rare hereditary problems of fructose intolerance should not take this medicine.

Taking STREPSILS STRAWBERRY LOZENGES with food and drink:

Interactions with food and drink for STREPSILS STRAWBERRY LOZENGES have not been established

Pregnancy and breast-feeding:

Safety in pregnancy and lactation has not been established

If you are pregnant or breast-feeding your baby, please consult your doctor, pharmacist or other health care professional for advice before taking this medicine **Driving and using machinery:**

No adverse effects known

Taking other medicines with Strepsils Strawberry lozenges:

Always tell your healthcare professional if you are taking any other medicine. (This includes complementary or traditional medicines).

4. HOW TO TAKE STREPSILS STRAWBERRY LOZENGES

The usual dose is:

Adults and children (6 years old and above):

One lozenge to be dissolved slowly in the mouth every 2-3 hours up to a maximum of 12 lozenges in 24 hours.

Elderly: there is no need for dosage reduction in the elderly.

Not suitable for children under 6 years.

Always take Strepsils Strawberry Lozenges exactly as instructed. You should check with your doctor or pharmacist if you are unsure.

Do not share medicines prescribed for you with any other person

If you take more STREPSILS STRAWBERYY LOZENGES than you should

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre

5. POSSIBLE SIDE EFFECTS

Strepsils Strawberry Lozenges can have side effects.

Not all side effects reported for Strepsils Strawberry Lozenges are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking this medicine, please consult your doctor, pharmacist or other healthcare professional for advice

If any of the following happens, stop taking Strepsils Strawberry Lozenges and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Swelling of the hands, feet, ankles, face, lips, mouth or throat, which may cause difficulty in swallowing or breathing.
- Rash or itching.

These are all very serious side effects. If you have them, you may have had a serious allergic reaction to Strepsils Strawberry Lozenges. You may need urgent medical attention or hospitalisation If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

6. STORING AND DISPOSING OF STREPSILS STRAWBERRY LOZENGES

Store all medicines out of reach of children

Store at or below 25°C

Do not use the product after the expiry date shown on the carton or blister

Return all unused medicine to your pharmacist

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets)

7. PRESENTATION

Cartons containing 16 lozenges in white opaque blister strips

8. IDENTIFICATION

A pink, circular lozenge with a characteristic strawberry taste and Strepsils brand icon intagliated on both sides

9. REGISTRATION NUMBER

A40/16.4/0358

10. NAME, BUSINESS ADDRESS AND TELEPHONE NUMBER OF THE HOLDEOF CERTIFICATE OF REGISTRATION

Reckitt Benckiser Pharmaceuticals (Pty) Ltd

8 Jet Park Road

Elandsfontein

1601

South Africa

+2711 871 1611

11. DATE OF PUBLICATION OF THIS PATIENT INFORMATION LEAFLET

Date on the registration certificate: 23 July 2010

Date of the most recently revised patient information leaflet: 13 December 2019

Please go to the link for access to the corresponding PI:

https://cpdcentre.co.za/productinserts/Strepsils-Strawberry.pdf

SKEDULERINGSTATUS

S0

EIENDOMSNAAM, STERKTE EN DOSEERVORM

STREPSILS STRAWBERRY SUIGTABLETTE. 2,4-dichloorbenzylalkohol 1,2 mg en amymetrakresol 0,6 mg.

Lees hierdie hele pamflet noukeurig deur want dit bevat belangrike inligting vir u.

Strepsils Strawberry Suigtablette is beskikbaar sonder ’n voorskrif van ’n dokter, vir u om matige siektes te behandel. Ongeag moet u steeds Strepsils Strawberry Suigtablette versigtig gebruik om die beste resultaat daaruit te kry.

- Bewaar hierdie pamflet. Dit mag nodig wees om dit weer te lees.
- Moet nie Strepsils Strawberry Suigtablette met enige ander persoon deel nie.
- Vra u apteker indien u meer inligting of advies benodig.
- U moet ’n dokter raadpleeg indien u simptome vererger of na 2 dae nie verbeter nie.

1. WAT STREPSILS STRAWBERRY SUIGTABLETTE BEVAT

- Die aktiewe stowwe is 2,4-dichloorbensielalkohol en amielmetakresol. Elke suigtablet bevat 2,4dichloorbenzylalkohol 1,2 mg en amylmetakresol 0,6 mg.
- Die ander bestanddele is Flav P Aarbei geursel 052312B, vloeistof, pienk antho P-WS (E163), gesuiwerde water en wynsteensuur.
- Bevat versoeter (isomalt 1830 mg, maltitol 481 mg, natriumsakkarien 2 mg per suigtablette)

2. WAARVOOR STREPSILS STRAWBERRY SUIGTABLETTE gebruik word

Strepsils Strawberry Suigtablette bevat antiseptiese bestanddele wat gebruik word vir die verligting van geringe keel- en mondinfeksies asook seer keel.

3. VOOR U STREPSILS STRAWBERRY SUIGTABLETTE NEEM

Moet nie Strepsils Strawberry Suigtablette:

Indien jy hipersensitief (allergies) is vir 2,4-dichloorbenzylalkohol, amielmetakresol, of vir enige van die ander bestanddele van Strepsils Strawberry Suigtablette.

Wees veral versigtig met STREPSILS STRAWBERRY SUIGTABLETTE

STREPSILS STRAWBERRY SUIGTABLETTE bevat isomaltitol en maltitolstroop, wat ’n ligte lakseermiddel effek kan hê as ’n hele aantal op ’n dag geneem word. Pasiënte met ’n raar oorerflike probleme van fruktose-onverdraagsaamheid moet nie hierdie medisyne gebruik nie.

Die neem van STREPSILS STRAWBERRY SUIGTABLETTE met spys en drank:

Geen interaksie met spys en drank vir STREPSILS STRAWBERRY SUIGTABLETTE is vasgestel nie.

Swangerskap en borsvoeding:

Veiligheid tydens swangerskap en laktasie is nog nie vasgestel nie.

Indien u swanger is of u baba borsvoed, raadpleeg u dokter, apteker of ander professionele gesondheidswerker vir advies voor die gebruik van hierdie medisyne.

Bestuur en gebruik van masjinerie:

Geen nadelige gevolge is bekend nie.

Die gebruik van ander medisyne saam met Strepsils Strawberry Suigtablette:

Lig altyd u professionele gesondheidswereker in as u enige ander medisyne neem. (insluitend aanvullende of tradisionele medisyne.)

4. HOE OM STREPSILS STRAWBERRY SUIGTABLETTE TE NEEM

Die gewone dosis is:

Volwassenes en kinders (6 jaar en ouer):

Een suigtablet moet elke 2-3 uur stadig in die mond opgelos word, tot en met ’n maksimum van 12 suitablette in 24 ure.

Bejaardes: die dosisvermindering by bejaardes is nie nodig nie Nie

geskik vir kinders jonger as 6 jaar oud nie.

Neem altyd Strepsils Strawberry Suigtablette presies soos voorgeskryf. Raadpleeg u dokter of apteker indien u onseker is.

Moet nie medisyne wat aan u voorgeskryf is, deel met enige ander persoon nie.

Indien jy meer STREPSILS STRAWBERRY SUIGTABLETTE inneem as wat jy behoort: In die geval van oordosering, raadpleeg u dokter of apteker. Indien daar nie een beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum.

5. MOONTLIKE NEWE-EFFEKTE

Strepsils Strawberry Suigtablette mag newe-effekte hê.

Nie alle newe-effekte wat vir Strepsils Strawberry Suigtablette aangemeld word, is in hierdie pamflet ingesluit nie. Indien u algemene gesondheid vererger of as u enige probleme ervaar tydens die gebruik van Strepsils Strawberry Suigtablette, raadpleeg u dokter, apteker of ander professionele gesondheidswerker vir advies.

Indien enige van die volgende plaasvind, hou op met die gebruik van Strepsils Strawberry Suigtablette en deel dit onmiddelik met u dokter of gaan na die ongevalle-afdeling van u naaste hospitaal:

- Swelling van die hande, voete, enkels, gesig, lippe, mond of keel, wat probleme met sluk of asemhaling kan veroorsaak.
- Uitslag of jeukerigheid.

Hierdie is alles baie ernstige newe-effekte. Indien dit plaasvind, het u moontlik ’n ernstige allergiese reaksie op Strepsils Strawberry Suigtablette gehad. U mag dringende mediese-hulp of hospitalisasie benodig.

Indien u let op enige newe-effekte wat nie hier genoem word nie, raadpleeg u dokter of apteker.

6. BERGINGSAANWYSINGS EN VERWYDERING VAN STREPSILS STRAWBERRY SUIGTABLETTE

Stoor alle medisyne buite die bereik van kinders.

Bewaar teen of benede 25˚C.

Moet nie die produk gebruik na verloop van die vervaldatum wat op die karton of blisterstrook aangedui word nie.

Besorg alle ongebruikte medisyne terug aan u apteker.

Moet nie wegdoen met ongebruikte medisyne in dreine en rioolstelsels (bv. Toilette) nie.

7. AANBIEDING

Karton bevat 16 suikertablette in wit ondeursigtige stulpstroke.

8. IDENTIFIKASIE

'n pienk, sirkelvormige suigtablet met 'n kenmerkende aarbei-smaak en die Strepsils-handelsmerk-ikoon wat aan beide kante uitgesky is.

9. REGISTRASIENOMMER

A40/16.4/0358

10. NAAM, BESIGHEIDSADRES EN TELEFOONNOMMER VAN DIE HOUER VAN DIE SERTIFIKAAT VAN REGISTRASIE

Reckitt Benckiser Pharmaceuticals (Edms) Bpk

8 Jet Park Road

Elandsfontein

1601

South Africa

+2711 871 1611

11. DATUM VAN PUBLIKASIE VAN DIE PASIËNT-INLIGTINGSVOUBILJET

Datum op die registrasiesertifikaat: 23 Julie 2010

Datum op die mees onlangste pasiënt-inligtingsvoubiljet: 13 Desember 2019

Gaan asseblief na die web adres vir die ooreenstemmende professionele informasie:

https://cpdcentre.co.za/product-inserts/Strepsils-Strawberry.pdf